

# $t$ -ANALOGUE OF $q$ -CHARACTERS AND BASES OF QUANTUM CLUSTER ALGEBRAS

FAN QIN

**ABSTRACT.** We introduce a new family of graded quiver varieties together with a new  $t$ -deformation of the associated Grothendieck rings. We generalize Nakajima's  $t$ -analogue of  $q$ -characters to these constructions. This allows a graded monoidal pseudo-categorification approach to bases of certain quantum cluster algebras. When the initial seed is acyclic, for any choice of coefficients and quantization, we obtain a generic basis, a dual PBW basis, and a dual canonical basis with positive structure constants. As a by-product, we obtain the quantum cluster characters for generic modules.

We conclude the paper by showing how a general quantum cluster algebra depends on the coefficients and the quantization. This provides a general framework for the correction techniques in our pseudo-categorification approach.

This paper provides the geometric foundations for a joint paper by Yoshiyuki Kimura and the author. When constructing the bases, we shall use some important results of the generic characters obtained in that paper via Fourier-Deligne-Sato transforms.

## CONTENTS

|                                                |    |
|------------------------------------------------|----|
| 1. Introduction                                | 1  |
| Acknowledgments                                | 5  |
| 2. Preliminaries                               | 5  |
| 3. Quantum cluster variables via normalization | 11 |
| 4. Graded quiver varieties                     | 12 |
| 5. $qt$ -characters                            | 24 |
| 6. Results on generic characters               | 32 |
| 7. From characters to the quantum torus        | 34 |
| 8. Bases of quantum cluster algebras           | 39 |
| 9. Change of coefficients and quantizations    | 45 |
| References                                     | 49 |

## 1. INTRODUCTION

**1.1. Motivation.** Cluster algebras were invented by Fomin and Zelevinsky in [FZ02]. These are algebras generated by certain combinatorially-defined generators (the cluster variables). The quantum deformations were defined in [BZ05]. Since its very beginning, the theory of (quantum) cluster algebras has been related to many areas, such as Poisson Geometry, discrete dynamical systems, higher Teichmüller spaces, combinatorics, commutative

and non-commutative algebraic geometries, and representation theory, cf. the introductory survey by Bernhard Keller [Kel12].

In this paper, we restrict our attentions to the (quantum) cluster algebras themselves. Fomin and Zelevinsky stated their original motivation as follows:

*This structure should serve as an algebraic framework for the study of dual canonical bases in these coordinate rings and their  $q$ -deformations. In particular, we conjecture that all monomials in the variables of any given cluster (the cluster monomials) belong to this dual canonical basis.*

Therefore, bases of cluster algebra deserve to be put under scrutiny. The following two questions arise naturally.

**Question 1.1.1.** *1) How to construct bases of (quantum) cluster algebras?  
2) What are the corresponding structure constants and transition matrices?*

**Question 1.1.2.** *1) Can we construct a “dual canonical basis” of the quantum cluster algebra, such that its structure constants are positive and it contains all the quantum cluster monomials.*

*2) Can we identify this basis with a subset of the dual canonical basis of certain quantum group?*

**Remark 1.1.3.** *We expect this basis of quantum cluster algebra to be identified with only a subset of the dual canonical basis, because the cluster algebras are “smaller” than general quantum groups, cf. [GLS11].*

Furthermore, by [FZ07] [Tra09], (quantum) cluster monomials vary little when the choice of the coefficient pattern changes. If such a result holds for the bases as well, our research in basis will be largely simplified. Surprisingly, at best of the author’s knowledge, this problem seems has not been studied in literature yet.

**Question 1.1.4.** *How does the bases data depend on the choice of the coefficients and quantization? Are they controllable?*

The final motivation of this paper comes from the study of quantum cluster characters. In [Qin10], the author defined quantum cluster characters over the rigid objects of certain cluster category, and showed that these characters describe the quantum cluster monomials of acyclic quantum cluster algebras. Nagao proposed a more general formula to describe the quantum cluster variables of general (quantum) cluster algebras based on the theory of non-commutative Donaldson-Thomas invariants developed by Kontsevich and Soibelman (up to some conjectures), cf. [KS08] [Nag10]. The following question is natural from the representation theoretic point of view.

**Question 1.1.5.** *Can we extend the quantum cluster character to generic objects, such that we obtain a generic basis of the quantum cluster algebra containing all the quantum cluster monomials?*

## 1.2. Previous context, strategies, and results.

*Previous context.* Despite the many successful applications of (quantum) cluster algebras to other areas, the basis construction problems which motivate their introduction remain largely open.

We have limited knowledge of the constructions of bases of the classical cluster algebras. In the approach via preprojective algebra, *cf.* [GLS11], Geiß, Leclerc, and Schroëer have shown that if  $G$  is a semi-simple complex algebraic group and  $N \subset G$  a maximal nilpotent subgroup, then the coordinate algebra  $\mathbb{C}[N]$  admits a canonical classical cluster structure whose coefficient type is specific. They further constructed the generic basis of  $\mathbb{C}[N]$ , which contain the cluster monomials, and identified it with Lusztig’s dual semicanonical basis of  $\mathbb{C}[N]$  [Lus00]. As another approach, recently, Musiker, Schiffler, and Williams constructed bases for classical cluster algebras arising from unpunctured surfaces with coefficients whose exchange matrix is of full rank, [MSW11].

Our knowledge of the bases of the quantum cluster algebras is even more limited. [Lam11a] [Lam11b] [DX11] [HL11] obtained partial results of Question 1.1.2 for quivers of finite and affine type. Standard bases and partial results of triangular bases have been obtained for acyclic quantum cluster, *cf.* [BFZ05] [BZ11] [BZ12].

*Deformed monoidal pseudo-categorification.* In this paper, we use (deformed) monoidal pseudo-categorifications to give a new approach to bases of (quantum) cluster algebras.

Monoidal categorification was used by Hernandez and Leclerc as a new approach to the positivity conjecture of cluster algebras, *cf.* [HL10]. For a given cluster algebra  $\mathcal{A}$ , we want to find a tensor category  $\mathcal{C}$  together with a well behaved algebra isomorphism from  $\mathcal{A}$  to the Grothendieck ring  $R$  of  $\mathcal{C}$ . In particular, each cluster monomial should be sent to the class of a simple module. In the original work [HL10], the tensor category  $\mathcal{C}$  is the tensor category of certain finite dimensional modules of certain quantum affine algebra, and the isomorphism is obtained by comparing the (truncated)  $q$ -characters of these modules with cluster characters.

Therefore, we easily arrive at the following naive idea: In order to constructing the bases of (quantum) cluster algebras: it suffices to study bases of the (deformed) Grothendieck ring, and then apply the algebra isomorphism.

In this paper, we use a similar category  $\mathcal{C}$  for quantum loop algebras and construct an linear map identifying the (truncated)  $qt$ -characters with quantum cluster characters studied by the author in [Qin10]. Our construction holds for general coefficients and quantizations, where this map fails to be algebraic, thus the name “pseudo-categorification”. A key ingredient is the observation that the failures are mild and controllable. Therefore, in practice, we can still follow the above naive idea to study bases.

*Graded quiver varieties.* In order to construct this monoidal pseudo-categorification, we need to understand the deformed Grothendieck ring and the  $tq$ -characters. A second key ingredient is that we can understand them geometrically by using Nakajima’s quiver varieties, *cf.* [Nak01a] [Nak04].

Recall that Nakajima’s quiver variety is just a natural generalization of the ADHM-construction of instantons. And his graded quiver variety is the fixed point subvariety with respect to a  $\mathbb{C}^*$ -action. In [Nak11], in constructing the graded quiver varieties, Nakajima required the quiver to be bipartite.

In Section 4, we introduce a new family graded quiver varieties for acyclic quivers which are not necessarily bipartite. In order to remove the bipartite restriction, we carefully change the definition of the  $\mathbb{C}^*$ -action, such that the resulting new graded quiver varieties (fixed point sets) still have good properties. Then we establish the geometric foundations of our discussion, and proceed to study the deformed Grothendieck rings and  $qt$ -characters following the arguments of Nakajima in [Nak01a] [Nak04] [Nak11].

Moreover, as an important application, these constructions lead to an affirmative answer of Question 1.1.2 for acyclic quantum cluster algebras of specific coefficients and quantization, which will be presented in the joint work [KQ12] by Yoshiyuki Kimura and the author. A corollary of the answer generalizes a result of Nakajima [Nak11]:

*When the quantum cluster algebra contains an acyclic seed, the positivity conjecture is true. Namely, the coefficients of the cluster expansions of the quantum cluster variables are positive with respect to any chosen seed.*

Its proof depends on the Fourier-Deligne-Sato transform, cf. the recent work by Efimov [Efi11] for an independent different approach. We shall use some useful results obtained in the proof.

*Main results.* Via the pseudo-categorification approach introduced above, we use  $t$ -analogue of  $q$ -characters to realize three bases of the acyclic quantum cluster algebras for any choice of coefficients and quantizations: a dual PBW basis, a canonical basis, and a generic basis. For such (quantum) cluster algebras, we obtain affirmative answers and solutions of Question 1.1.1, 1.1.2(1), 1.1.5. We also use the correction techniques to give an affirmative answer to Question 1.1.4, which generalizes the results in [FZ07] from cluster variables to bases.

**Remark 1.2.1.** *Notice that, by the joint work of Kimura with the author [KQ12], 1.1.2(2)] has an affirmative answer for quantum cluster algebra of a specific choice of coefficients and quantization, while for a general choice, a reasonable identification which preserves the multiplications does not exist.*

It should be mentioned that, despite that the results on the generic quantum cluster characters and the quantum generic basis seem very natural, their proof needs the full power of the graded quiver varieties and the existence of the monoidal categorification. For the moment, the author does not see any alternative approach.

**1.3. Plan of the paper.** In the preliminary section 2, we recall the definition of quantum cluster algebras arising from ice quivers together with their categorifications.

In section 3, we show how the cluster expansions of quantum cluster variables (monomials) of quantum cluster algebras with arbitrary compatible pairs can be deduced from those with unitally compatible pairs.

In section 1.2, we introduce a family of new graded quiver varieties by choosing a new torus action. Then we follow the statements of [Nak11] to construct the deformed Grothendieck ring  $R_t$ . In section 5, we define the  $t$ -analogue of  $q$ -characters for these graded quiver varieties.

In section 7, we measure the failure of these characters being isomorphism.

Our constructions allow to quantize [Nak11] by comparing the  $qt$ -characters with the quantum cluster characters in [Qin10]. This step is of independent interest and importance, and the proofs are contained in a separated paper [KQ12].

In Section 6, we recall some results on the generic characters obtained in [KQ12]. With this results, in Section 8, we obtain bases of acyclic quantum cluster algebra for any choice of coefficients and quantizations together with their structure constants and transition matrices.

Finally, in the last section, we prove the correction techniques which tells us how a general quantum cluster algebra depends on the choice of coefficients and quantization.

## ACKNOWLEDGMENTS

The author would like to express his sincere thanks to his thesis advisor Bernhard Keller for the encouragement and discussions. He also thank Yoshiyuki Kimura for many useful discussions, in particular for the explanations of his previous work on the factorization of dual canonical basis and of Nakajima's quiver varieties.

## 2. PRELIMINARIES

**2.1. Quantum cluster algebras.** We refer the reader to [Qin10] for detailed definitions and important properties.

Following [BZ05], we define (generalized) quantum cluster algebras over  $(R, v)$ , where  $R$  is an integral domain and  $v$  an invertible element in  $R$ . In the present paper we shall only be interested in the case where  $(R, v) = (\mathbb{Z}[v^\pm], v)$  for a formal parameter  $v$ .

Let  $m \geq n$  be two positive integers. Let  $\Lambda$  be an  $m \times m$  skew-symmetric integer matrix and  $\tilde{B}$  an  $m \times n$  integer matrix. The upper  $n \times n$  submatrix of  $\tilde{B}$ , denoted by  $B$ , is called the *principal part* of  $\tilde{B}$ .

**Definition 2.1.1** (Compatible pair). *The pair  $(\Lambda, \tilde{B})$  is called compatible if we have*

$$(1) \quad \Lambda(-\tilde{B}) = \begin{bmatrix} D \\ 0 \end{bmatrix}$$

*for some  $n \times n$  diagonal matrix  $D$  whose diagonal entries are strictly positive integers. It is called a unitally compatible pair if moreover  $D$  is the identity matrix  $\mathbf{1}_n$ .*

Let  $(\Lambda, \tilde{B})$  be a compatible pair. The component  $\Lambda$  is called the  $\Lambda$ -matrix of  $(\Lambda, \tilde{B})$ , and the component  $\tilde{B}$  the  $B$ -matrix of  $(\Lambda, \tilde{B})$ .

**Proposition 2.1.2.** [BZ05, Proposition 3.3] *The  $B$ -matrix  $\tilde{B}$  has full rank  $n$ , and the product  $DB$  is skew-symmetric.*

We write  $\Lambda(g, h)$  for  $g^T \Lambda h$ ,  $g, h \in \mathbb{Z}^m$ , where  $(\ )^T$  means taking the matrix transposition.

**Definition 2.1.3** (Quantum torus). *The quantum torus  $\mathcal{T} = \mathcal{T}(\Lambda)$  over  $(R, v)$  is the Laurent polynomial ring  $\mathbb{Z}[v^\pm][x_1^\pm, \dots, x_m^\pm]$ , endowed with the twisted product  $*$  such that we have*

$$x^g * x^h = v^{\Lambda(g, h)} x^{g+h}$$

for any  $g$  and  $h$  in  $\mathbb{Z}^m$ . Here for any  $g = (g_i)_{1 \leq i \leq m} \in \mathbb{Z}^m$ ,  $x^g$  denote the monomial  $\prod_{1 \leq i \leq m} X_i^{g_i}$ .

We denote the usual product in  $\mathcal{T}$  by  $\cdot$ , and often omit this notation.

Assume that  $R$  is endowed with an involution sending each element  $r$  to  $\bar{r}$ , such that  $\bar{\bar{v}}$  equals  $v^{-1}$ . We extend the involution of  $R$  to an involution (anti-automorphism) of  $\mathcal{T}$  by defining  $\overline{x^g} = x^g$  for any  $g \in \mathbb{Z}^m$ .

A sign  $\epsilon$  is an element in  $\{-1, +1\}$ . Denote by  $b_{ij}$  the entry in position  $(i, j)$  of  $\tilde{B}$ . For any  $1 \leq k \leq n$  and any sign  $\epsilon$ , we associate to  $\tilde{B}$  an  $m \times m$  matrix  $E_\epsilon$  whose entry in position  $(i, j)$  is

$$(2) \quad e_{ij} = \begin{cases} \delta_{ij} & \text{if } j \neq k \\ -1 & \text{if } i = j = k \\ \max(0, -\epsilon b_{ik}) & \text{if } i \neq k, j = k, \end{cases}$$

and an  $n \times n$  matrix  $F_\epsilon$  whose entry in position  $(i, j)$  is

$$f_{ij} = \begin{cases} \delta_{ij} & \text{if } i \neq k \\ -1 & \text{if } i = j = k \\ \max(0, \epsilon b_{kj}) & \text{if } i = k, j \neq k. \end{cases}$$

Fix a compatible pair  $(\Lambda, \tilde{B})$  and the quantum torus  $\mathcal{T} = \mathcal{T}(\mathbb{Z}^m, \Lambda)$ . Notice that the quantum torus  $\mathcal{T} = \mathcal{T}(+, *)$  is contained in its skew-field of fractions, which is denoted by  $\mathcal{F}$ , cf. [BZ05, Appendix].

In the following, we consider triples  $(\Lambda', \tilde{B}', x')$  such that  $(\Lambda', \tilde{B}')$  is a compatible pair and  $x' = (x'_1, \dots, x'_m)$  is an  $m$ -tuple of elements in the quantum torus  $\mathcal{F}$ .

Let  $\mathbb{T}_n$  be an  $n$ -regular tree with root  $t_0$ . There is a unique way of associating a triple  $(\Lambda(t), \tilde{B}(t), x(t))$  with each vertex  $t$  of  $\mathbb{T}_n$  such that we have

- (1)  $(\Lambda(t_0), \tilde{B}(t_0), x(t_0)) = (\Lambda, \tilde{B}, x)$ , where  $x = (x_1, \dots, x_m)$ , and
- (2) if two vertices  $t$  and  $t'$  are linked by an edge labeled  $k$ , then the seed  $(\Lambda(t'), \tilde{B}(t'), x(t'))$  is obtained from  $(\Lambda(t), \tilde{B}(t), x(t))$  by the mutation at  $k$  defined as below.

**Definition 2.1.4** (Mutation [BZ05]). *Given any sign  $\epsilon$ , the new triple  $(\Lambda(t'), \tilde{B}(t'), x(t'))$  obtained from  $(\Lambda(t), \tilde{B}(t), x(t))$  by the mutation at  $k$  is given by*

$$(3) \quad (\Lambda(t'), \tilde{B}(t')) = (E_\epsilon(t)^T \Lambda(t) E_\epsilon(t), E_\epsilon(t) \tilde{B}(t) F_\epsilon(t)),$$

and

$$(4) \quad \begin{aligned} x_k(t) * x_k(t') &= v^{\Lambda(t)(e_k, \sum_{1 \leq i \leq m} [b_{ik}(t)] + e_i)} \prod_{1 \leq i \leq m} x_i(t)^{[b_{ik}] + (t)} \\ &+ v^{\Lambda(t)(e_k, \sum_{1 \leq j \leq m} [-b_{jk}(t)] + e_j)} \prod_{1 \leq j \leq m} x_j(t)^{[-b_{jk}] + (t)}, \end{aligned}$$

$$(5) \quad x_i(t') = x_i(t), \quad 1 \leq i \leq m, \quad i \neq k.$$

We recall from [BZ05] that  $\mu_k$  is an involution, and is independent of the choice of  $\epsilon$ .

The triples  $(\Lambda(t), \tilde{B}(t), x(t))$ ,  $t \in \mathbb{T}_n$ , are called the (quantum) seeds. The elements  $x_i(t)$ ,  $1 \leq i \leq m$ ,  $t \in \mathbb{T}_n$ , are called the  $x$ -variable. The  $x$ -variables  $x_i(t)$ ,  $1 \leq i \leq n$ ,  $t \in \mathbb{T}_n$ , are called the *quantum cluster variables*, and the monomials (with respect to the usual product  $\cdot$ ) in them are called the *quantum cluster monomials*. Notice that for  $j > n$ , the  $x$ -variables  $x_j(t)$  do not depend on  $t$ .

**Definition 2.1.5** (Quantum cluster algebra). *The quantum cluster algebra  $\mathcal{A}^q = \mathcal{A}^q(+, *)$  over  $(R, v)$  is the  $R$ -subalgebra of  $\mathcal{F}$  generated by the quantum cluster variables  $x_i(t)$  for all the vertices  $t$  of  $\mathbb{T}_n$  and  $1 \leq i \leq n$ , and the elements  $x_j$  and  $x_j^{-1}$  for all  $j > n$ .*

The specialization  $\mathcal{A} = \mathcal{A}^q|_{v \mapsto 1}$  is called the *classical cluster algebra*, which is also denoted by  $\mathcal{A}^{\mathbb{Z}}$ . If  $R = \mathbb{Z}[v^{\pm}]$ , we say that  $\mathcal{A}^q$  and  $\mathcal{A}$  are *integral*.

**Theorem 2.1.6** (Quantum Laurent phenomenon). [BZ05, Section 5] *The quantum cluster algebra  $\mathcal{A}^q$  is a subalgebra of  $\mathcal{T}$ .*

Similarly, the cluster algebra  $\mathcal{A} = \mathcal{A}^{\mathbb{Z}}$  is a subalgebra of the Laurent polynomial ring  $\mathcal{T}|_{v \mapsto 1} = \mathcal{T}^{\mathbb{Z}}$ .

**2.2. Coefficient types (frozen patterns).** A quiver  $Q$  is an oriented graph, which consists of a set of vertices  $I = \{1, \dots, n\}$  and a set of arrows  $\Omega$ . For each arrow  $h$ , its source is denoted by  $s(h)$  and its target by  $t(h)$ . Associate to  $h$  a new arrow  $\bar{h}$  which points from  $t(h)$  to  $s(h)$ . Denote the set  $\{\bar{h} | h \in \Omega\}$  by  $\bar{\Omega}$ . Define  $H$  to be the disjoint union of  $\Omega$  and  $\bar{\Omega}$ .  $Q$  is called acyclic if it contains no oriented cycles. The opposite quiver  $Q^{op}$  of  $Q$  consists of the vertices in  $I$  and the arrows in  $\bar{\Omega}$ . Sometimes we also denote  $I$  and  $\Omega$  by  $Q_0$  and  $Q_1$  respectively.

**Example 2.2.1.** *The acyclic quiver  $Q$  in Figure 1 has vertices 1, 2, 3. Its opposite quiver  $Q^{op}$  is given by Figure 2.*






FIGURE 1. An acyclic quiver  $Q$

Let  $Q$  be a full subquiver of another quiver  $\tilde{Q}$ , which has vertices  $\{1, \dots, m\}$  and a set of arrows  $\tilde{\Omega}$ . We say  $\tilde{Q}$  is an ice quiver with principal part  $Q$  and coefficient type (or frozen pattern)  $\tilde{\Omega} - \Omega$ . The vertices  $n + 1, \dots, m$  are called *frozen vertices*.

FIGURE 2. The quiver  $Q^{op}$ 

**Example 2.2.2.** Figure 3 is an example of an ice quiver with  $m = 6$ , whose principal part is the quiver in Figure 1.

FIGURE 3. An ice quiver  $\tilde{Q}_1^z$  of level 1 with  $z$ -pattern

We associate to  $\tilde{Q}$  an  $m \times n$  matrix<sup>1</sup>  $\tilde{B} = (b_{ij})$ , whose coefficient  $b_{ij}$  is the difference between the number of arrows from  $i$  to  $j$  and the number of arrows from  $j$  to  $i$  in  $\tilde{Q}$ . Similarly, we associate to  $Q$  an  $n \times n$ -matrix  $B = B_Q$ . If further a compatible pair  $(\Lambda, \tilde{B})$  is given, we can construct the associated quantum cluster algebra  $\mathcal{A}^q$ .

**Example 2.2.3.** The matrix  $B = B_Q$  associated with the quiver  $Q$  in Figure 1 is

$$\begin{pmatrix} 0 & 1 & 1 \\ -1 & 0 & 1 \\ -1 & -1 & 0 \end{pmatrix}.$$

The matrix  $\tilde{B}$  associated to the ice quiver  $\tilde{Q}$  in Figure 3 is

$$\begin{pmatrix} 0 & 1 & 1 \\ -1 & 0 & 1 \\ -1 & -1 & 0 \\ -1 & 0 & 0 \\ 1 & -1 & 0 \\ 1 & 1 & -1 \end{pmatrix}.$$

<sup>1</sup>Notice that this convention is opposite to that of [Nak11].



Denote the quotient functor by  $\pi : \text{per } \Gamma \rightarrow \mathcal{C}$  and define

$$\begin{aligned} T_i &= \pi(e_i \Gamma), \quad 1 \leq i \leq m \\ T &= \bigoplus_{1 \leq i \leq m} T_i. \end{aligned}$$

It is shown in [Pla11] that the endomorphism algebra of  $T$  is isomorphic to  $H^0 \Gamma$ .

For any triangulated category  $\mathcal{U}$  and any rigid object  $X$  of  $\mathcal{U}$ , we define the subcategory  $\text{pr}_{\mathcal{U}}(X)$  of  $\mathcal{U}$  to be the full subcategory consisting of the objects  $M$  such that there exists a triangle in  $\mathcal{U}$

$$M_1 \rightarrow M_0 \rightarrow M \rightarrow \Sigma M_1,$$

for some  $M_1$  and  $M_0$  in  $\text{add } X$ . The *presentable cluster category*  $\mathcal{D} \subset \mathcal{C}$  is defined as the full subcategory consisting of the objects  $M$  such that

$$M \in \text{pr}_{\mathcal{C}}(T) \cap \text{pr}_{\mathcal{C}}(\Sigma^{-1}T) \quad \text{and} \quad \dim \text{Ext}_{\mathcal{C}}^1(T, M) < \infty,$$

cf. [Pla10b].

We refer to [Pla10b] for the definition of the iterated mutations of the object  $T$ . There is a unique way of associating an object  $T(t) = \bigoplus_{1 \leq i \leq m} T_i(t)$  of  $\mathcal{D}$  with each vertex  $t$  of  $\mathbb{T}_n$  such that we have

- (1)  $T(t_0) = T$ , and
- (2) if two vertices  $t$  and  $t'$  are linked by an edge labeled  $k$ , then the object  $T(t')$  is obtained from  $T(t)$  by the mutation at  $k$ .

Let  $\mathcal{F} \subset \text{per } \Gamma$  denote the full subcategory  $\text{pr}_{\text{per } \Gamma}(\Gamma)$ . The quotient functor  $\pi : \text{per } \Gamma \rightarrow \mathcal{C}$  induces an equivalence  $\mathcal{F} \xrightarrow{\sim} \text{pr}_{\mathcal{C}}(T)$ . Denote by  $\pi^{-1}$  the inverse equivalence. For an object  $M \in \text{pr}_{\mathcal{C}}(T)$ , we define its *index*  $\text{ind}_T M$  as the class  $[\pi^{-1}M]$  in  $\text{K}_0(\text{per } \Gamma)$ .

**Theorem 2.3.1.** [Pla10b] (1) For any vertex  $t$  of  $\mathbb{T}_n$ , the classes  $[\text{ind}_T T_i(t)]$  form a basis of  $\text{K}_0(\text{per } \Gamma)$ .

(2) For a class  $[P]$  in  $\text{K}_0(\text{per } \Gamma)$ , let  $[[P] : T_i(t)]$  denote its  $i$ th coordinate in this basis. Then we have  $([\text{ind}_T T_i(t) : T_j])_{1 \leq j \leq m} = \tilde{g}_i(t)$ .

**Definition 2.3.2** (Coefficient-free objects). An object  $M$  in  $\mathcal{C}$  is called coefficient-free if

- (1) the object  $M$  does not contain a direct summand  $T_i$ ,  $i > n$ , and
- (2) the space  $\text{Ext}_{\mathcal{C}}^1(T_i, M)$  vanishes for  $i > n$ .

For a coefficient-free object  $M \in \mathcal{C}$ , the space  $\text{Ext}_{\mathcal{C}}^1(T, M)$  is a right  $H^0 \Gamma$ -module whose support is concentrated on  $Q$ . Thus, it can be viewed as a  $\mathcal{P}(Q, W)$ -module, where  $W$  is the potential on  $Q$  obtained from  $\tilde{W}$  by deleting all cycles through vertices  $j > n$  and  $\mathcal{P}(Q, W)$  is the Jacobi algebra of  $(Q, W)$ . Denote by  $\phi : \text{K}_0(\text{mod } \mathcal{P}(Q, W)) \rightarrow \text{K}_0(\text{per } \Gamma)$  the map induced by the composition of inclusions  $\text{mod } \mathcal{P}(Q, W) \rightarrow \mathcal{D}_{fd} \Gamma \rightarrow \text{per } \Gamma$ . For any vertex  $i$  of  $\tilde{Q}$ , it sends  $[S_i]$  to

$$(6) \quad \sum_{\text{arrows } i \rightarrow j} [e_j \Gamma] - \sum_{\text{arrows } l \rightarrow i} [e_l \Gamma],$$

as one easily checks using the minimal cofibrant resolution of the simple dg  $\Gamma$ -module  $S_i$ , cf. [KY11].

By the *twisted Poincaré polynomial* of a topological space  $Z$ , we mean the polynomial  $p_t(Z) = \sum_p (-1)^p \dim H^p(Z, \mathbb{Q})$ . When  $Q$  is acyclic, we have the following construction.

**Definition 2.3.3** (Quantum CC-formula, [Qin10]). *For any coefficient-free and rigid object  $M \in \mathcal{D}$ , we denote by  $m$  the class of  $\text{Ext}_{\mathcal{C}}^1(T, M)$  in  $K_0(\text{mod } kQ)$ , and associate to  $M$  the following element in  $\mathcal{T}^{\mathbb{Z}}$ :*

$$(7) \quad x_M = \sum_e p_{q^{\frac{1}{2}}}(\text{Gr}_e(\text{Ext}_{\mathcal{C}}^1(T, M))) q^{-\frac{1}{2} \dim \text{Gr}_e(\text{Ext}_{\mathcal{C}}^1(T, M))} x^{\text{ind}_T(M) - \phi(e)},$$

where  $p_{q^{\frac{1}{2}}}(\ )$  denote the twisted Poincaré polynomial, and  $\text{Gr}_e(\text{Ext}_{\mathcal{C}}^1(T, M))$  is the submodule Grassmannian of  $\text{Ext}_{\mathcal{C}}^1(T, M)$  whose  $\mathbb{C}$ -points are the submodules of the class  $e$  in  $K_0(\text{mod } \mathcal{P}(Q, W))$ .

The following theorem is the main result of [Qin10].

**Theorem 2.3.4** ([Qin10]). *Assume that the quiver  $Q$  is acyclic. For any vertex  $t$  of  $\mathbb{T}_n$  and any  $1 \leq i \leq n$ , we have*

$$x_{T_i(t)} = x_i(t).$$

Moreover, the map taking an object  $M$  to  $X_M$  induces a bijection from the set of isomorphism classes of coefficient-free rigid objects of  $\mathcal{D}$  to the set of quantum cluster monomials of  $\mathcal{A}^q$ .

### 3. QUANTUM CLUSTER VARIABLES VIA NORMALIZATION

Let  $\tilde{Q}$  be an ice quiver with principal part  $Q$  and  $\tilde{B}$  its associated  $m \times n$  matrix. Let  $\mathcal{A}^q$  be the quantum cluster algebra associated with a compatible pair  $(\Lambda, \tilde{B})$  such that  $\Lambda(-\tilde{B}) = \begin{bmatrix} D \\ 0 \end{bmatrix}$ .

Following [Pla10b], let  $\mathcal{C}_{\tilde{Q}}$  be the cluster category associated with the ice quiver  $\tilde{Q}$ . Then to each  $i \in I$  and  $t \in \mathcal{T}_n$ , associate an object  $M_i(t)$  as in Section 2.3 (where  $M_i(t)$  is denoted by  $T_i(t)$ ) such that  $M_i(t_0)[-1]$  is the projective module  $P_i$  via the canonical embedding of  $\text{mod } -\mathbb{C}Q$  into  $\mathcal{C}_{\tilde{Q}}$ . Furthermore, we have a map  $\text{ind}(\ )$  sending each object in  $\mathcal{C}_{\tilde{Q}}$  to vector  $\text{ind}(M_i(t))$  in  $\mathbb{Z}^m$ .

First choose  $\tilde{Q} - Q$  and  $\Lambda$  such that  $D$  is the identity matrix  $\mathbf{1}_n$ . The quantum cluster variables take the form

$$(8) \quad x_i(t) = \sum_{v \in \mathbb{N}^n} c_{i,v}(q^{\frac{1}{2}}) x^{\text{ind}(M_i(t)) + \tilde{B}v},$$

for some bar-invariant Laurent polynomials  $c_{i,v} \in \mathbb{Z}[t^{\pm}]$ . For general  $\tilde{Q} - Q$  and  $\Lambda$ , we have the following result.

**Theorem 3.0.5.** *Assume  $D$  is a matrix whose diagonal entries are equal to  $\delta \in \mathbb{Z}_{>0}$ . Then in  $\mathcal{A}^q$ , we have*

$$(9) \quad x_i(t) = \sum_{v \in \mathbb{N}^n} c_{i,v}(q^{\frac{\delta}{2}}) x^{\text{ind}(M_i(t)) + \tilde{B}v}.$$

*Proof.* Consider the  $2n \times 2n$ -matrix

$$\begin{pmatrix} B & -\mathbf{1}_n \\ \mathbf{1}_n & 0 \end{pmatrix}.$$

Denote its inverse by  $-\Lambda_1$ . Its submatrix  $\tilde{B}' = \begin{pmatrix} B \\ \mathbf{1}_n \end{pmatrix}$  is the  $2n \times n$ -matrix associated with the *ice quiver with principal coefficient*  $\tilde{Q}'$  and principal part  $Q$ . We have the objects  $M'_i(t)$  in the cluster category  $\mathcal{C}_{\tilde{Q}'}$ , and the map  $\text{ind}'$ .

Notice that  $(\Lambda_1, \tilde{B}')$  is a unitally compatible pair, and we have the associated quantum cluster algebra  $\mathcal{A}^{q_1}$ .

Inspired by [Tra09, Proposition 5.1], we consider the  $2n \times 2n$ -matrix

$$\Lambda_2 = \begin{pmatrix} 0 & -D \\ D & BD \end{pmatrix}.$$

The pair  $(\Lambda_2, \tilde{B}')$  is again compatible. In fact, one can check that  $\Lambda_2 = \delta \cdot \Lambda_1$ . Denote the associated quantum cluster algebra by  $\mathcal{A}^{q_2}$ .

Since  $\Lambda_2$  is divisible by  $\delta$ , we can view  $\mathcal{A}^{q_2}$  as a quantum cluster algebra over  $(\mathbb{Z}[v^\pm], v)$  with  $v = q^{\frac{\delta}{2}}$  and the initial compatible pair is  $(\Lambda_2/\delta, \tilde{B}')$ . Then by sending  $q^{\frac{1}{2}}$  to  $q^{\frac{\delta}{2}}$ ,  $\mathcal{A}^{q_1}$  is identified with  $\mathcal{A}^{q_2}$ . In  $\mathcal{A}^{q_2}$ , we have

$$\begin{aligned} x_i(t) &= \sum_{v \in \mathbb{N}^n} c_{i,v}(q^{\frac{\delta}{2}}) x^{\text{ind}'(M'_i(t)) + \tilde{B}'v} \\ &= \left( \sum_v c_{i,v}(q^{\frac{\delta}{2}}) q^{-\frac{1}{2}\Lambda_2(\tilde{B}'v, \tilde{g}_{i,1})} x^{\tilde{B}'v} \right) * x^{\text{ind}'(M'_i(t))} \end{aligned}$$

for the vectors  $\tilde{g}_{i,1} = \text{ind}'(M'_i(t))$ .

Finally, by [Tra09, Theorem 5.3] quantum  $F$ -polynomials exist. Therefore, the quantum cluster variables in  $\mathcal{A}^q$  can be written as

$$\begin{aligned} x_i(t) &= \left( \sum_v c_{i,v}(q^{\frac{\delta}{2}}) q^{-\frac{1}{2}\Lambda_2(\tilde{B}'v, \tilde{g}_{i,1})} x^{\tilde{B}'v} \right) * x^{\text{ind}(M_i(t))} \\ &= \sum_{v \in \mathbb{N}^n} q^{\frac{1}{2}f_{i,v}} c_{i,v}(q^{\frac{\delta}{2}}) x^{\text{ind}(M_i(t)) + \tilde{B}v}. \end{aligned}$$

for some integers  $f_{i,v} \in \mathbb{Z}$ . Notice that  $\{X^{\tilde{g}_i + \tilde{B}v} | v \in \mathbb{N}^n\}$  is linearly independent over  $\mathbb{Z}[q^{\pm\frac{1}{2}}]$  because we can show that  $\tilde{B}$  is full rank. Since  $X_i(t)$  and  $c_{i,v}$  are bar-invariant, all  $f_{i,v}$  must vanish.  $\square$

#### 4. GRADED QUIVER VARIETIES

We begin with constructing some graded quiver varieties associated with acyclic quivers. Our graded quiver varieties are different from the original ones in [Nak01a]. Thanks to our modified definition, we can show the existence of suitable stratifications for arbitrary acyclic quivers (Proposition 4.3.5). To prove this and other geometric properties, we adapt Nakajima's original arguments in [Nak98], [Nak01a]. For the convenience of the reader, we shall briefly review some important techniques, results, and give brief proofs of some properties.

Notice that, by the dimension of a complex variety, we always mean the complex dimension. By default, we *only* consider geometric points.

**Remark 4.0.6.** *It seems likely that our definition should still enjoy the link with quantum loop algebras established in [Nak01a]. However, since Lie theory is not the major focus in this thesis, we do not recheck the link. Therefore, when we follow any useful arguments in [Nak01a], the Lie theoretic expressions should be replaced with their geometric counterparts.*

**4.1. Graded quiver varieties.** Assume that  $Q$  is an acyclic quiver with the set of vertices  $I = \{1, \dots, n\}$ , such that  $b_{ij} \leq 0$  whenever  $i \geq j$ , cf. Section 2.2 for the definition of  $b_{ij}$ .

We denote the finitely supported bigraded vectors in  $\mathbb{N}^{I \times \mathbb{Z}}$  by  $w = (w_i(a))_{i \in I, a \in \mathbb{Z}}$ , and the finitely supported bigraded vectors in  $\mathbb{N}^{I \times (\mathbb{Z} + \frac{1}{2})}$  by  $v = (v_i(a))_{i \in I, a \in \mathbb{Z} + \frac{1}{2}}$ . Denote the associated bigraded complex vector spaces by  $W = \mathbb{C}^w = \bigoplus_{i,a} W_i(a) = \bigoplus_{i,a} \mathbb{C}^{w_i(a)}$  and similarly  $V = \mathbb{C}^v = \bigoplus_{i,a} V_i(a) = \bigoplus_{i,a} \mathbb{C}^{v_i(a)}$ .

The vectors  $w$  and  $v$  can be naturally viewed as elements in  $\mathbb{Z}^{I \times \mathbb{R}}$ . For any  $d \in \mathbb{R}$ , define the degree shift  $[d]$  of vectors  $\eta = \eta_i(a) \in \mathbb{Z}^{I \times \mathbb{R}}$  by  $\eta[d]_i(a) = \eta_i(a + d)$ . For any two vectors  $\eta^1 = \eta_i^1(a)$ ,  $\eta^2 = \eta_i^2(a)$  in  $\mathbb{Z}^{I \times \mathbb{R}}$ , if at least one of them has finite support, their inner product is defined as  $\eta^1 \cdot \eta^2 = \sum_{i,a} \eta_i^1(a) \eta_i^2(a)$ .

The Cartan matrix  $C$  associated with  $Q$  is the  $I \times I$  matrix whose entry in position  $(i, j)$  is

$$(10) \quad c_{ij} = \begin{cases} 2 & \text{if } i = j \\ -|b_{ij}| & \text{if } i \neq j \end{cases}.$$

**Definition 4.1.1** ( $q$ -Cartan matrix). *We define the linear map*

$$C_q : \mathbb{Z}^{I \times (\frac{1}{2} + \mathbb{Z})} \rightarrow \mathbb{Z}^{I \times \mathbb{Z}}$$

*such that for each  $\eta \in \mathbb{Z}^{I \times (\frac{1}{2} + \mathbb{Z})}$ , we have*

$$(11) \quad (C_q \eta)_k(a) = \eta_k(a + \frac{1}{2}) + \eta_k(a - \frac{1}{2}) - \sum_{i: 1 \leq i < k} b_{ik} \eta_i(a + \frac{1}{2}) - \sum_{j: k < j \leq n} b_{kj} \eta_j(a - \frac{1}{2}).$$

*It is called a  $q$ -analogue of the Cartan matrix  $C$ , or a  $q$ -Cartan matrix for short.*

It is easy to see that  $C_q$  naturally extends to a linear map from  $\mathbb{Z}^{I \times \mathbb{R}}$  to  $\mathbb{Z}^{I \times \mathbb{R}}$ , which we also denote by  $C_q$ . Furthermore,  $C_q$  commutes with  $[d]$ ,  $d \in \mathbb{R}$ .

The  $q$ -Cartan matrix  $C_q$  is symmetric in the following sense.

**Lemma 4.1.2.** *For any two vectors  $\eta^1, \eta^2 \in \mathbb{Z}^{I \times \mathbb{R}}$ , if at least one of them has finite support, we have the following identity:*

$$(12) \quad \eta^2 \cdot C_q \eta^1[-\frac{1}{2}] = C_q \eta^2 \cdot \eta^1[-\frac{1}{2}].$$

*Proof.* To prove the equation, it suffices to expand both sides:

$$\begin{aligned}
\text{LHS} &= \sum_{k,a} \eta_k^2(a) \cdot (\eta_k^1(a) + \eta_k^1(a-1) - \sum_{i:i < k} b_{ik} \eta_i^1(a) - \sum_{j:k < j} b_{kj} \eta_j^1(a-1)) \\
&= \sum_{k,a} (\eta_k^2(a - \frac{1}{2}) + \eta_k^2(a + \frac{1}{2}) - \sum_{j:k < j} b_{kj} \eta_j^2(a - \frac{1}{2}) \\
&\quad - \sum_{i:i < k} b_{ik} \eta_i^2(a + \frac{1}{2})) \cdot \eta_k^1(a - \frac{1}{2}) \\
&= \text{RHS}.
\end{aligned}$$

□

**Definition 4.1.3** (*l-Dominance*). A pair  $(v, w)$  is called *l-dominant*<sup>3</sup> if  $w - C_q v \in \mathbb{N}^{I \times \mathbb{Z}}$ .

Define the dominance order on the set of pairs  $(v, w)$ , such that  $(v', w') \leq (v, w)$  if  $w' - C_q v' = w - C_q(v + v'')$ , for some  $v'' \in \mathbb{N}^{I \times (\mathbb{Z} + \frac{1}{2})}$ .

We say  $v' \leq v$  if  $(v', w) \leq (v, w)$ , or equivalently, if  $v'_i(a) \geq v_i(a)$ , for all  $(i, a)$ . We say  $w' \leq w$  if  $(0, w') \leq (0, w)$ .

**Definition 4.1.4** (*Weight order*). We define the total order (weight order)  $<_w$  on the set of pairs  $(i, a)$ ,  $i \in I$ ,  $a \in \mathbb{R}$ , such that  $(i', a') <_w (i, a)$  if  $a' < a$ , or if  $a' = a$  and  $i' < i$ .

Denote the set  $\{\eta \in \mathbb{Z}^{I \times \mathbb{Z}} \mid \eta_k(a) = 0, \forall k \in I, a \ll 0\}$  by  $E$ . Denote the set  $\{\eta[\frac{1}{2}] \mid \eta \in E\}$  by  $E[\frac{1}{2}]$ .

**Lemma 4.1.5.** 1)  $C_q$  restricts to an isomorphism from  $E[\frac{1}{2}]$  to  $E$ .

2)  $C_q$  restricts to an isomorphism from  $E$  to  $E[-\frac{1}{2}]$ .

*Proof.* Since  $C_q$  commutes with  $[\frac{1}{2}]$ , it suffices to show 1). Consider the block matrix of the restricted map  $C_q[\frac{1}{2}]$  from  $E$  to  $E$ , whose components are indexed by  $(i, a) \times (j, b)$ ,  $i, j \in I$ ,  $a, b \in \mathbb{Z}$ . It is lower unitriangular with respect to the weight order. Notice that  $E$  is bounded below. The claim follows. □

We denote the inverses of both the restriction maps by  $C_q^{-1}$ .

Let  $e_{k,a} \in \mathbb{N}^{I \times \mathbb{Z}}$  be the unit vector concentrated at the degree  $(k, a)$ . Denote by  $p_{i,j}$  the dimension of  $\text{Hom}_{\text{mod } \mathbb{C}Q}(P_i, P_j)$ ,  $i \leq j$ , namely the number of (possibly trivial) paths from  $i$  to  $j$  in the quiver  $Q$ .

**Lemma 4.1.6.** The vector  $\tilde{e}_{k,a} = C_q^{-1}(e_{k,a})$  satisfies, for all  $k' \in I$ ,

$$(13) \quad \tilde{e}_{k,a} \cdot e_{k',b} = \begin{cases} 0 & \text{if } b < a + \frac{1}{2} \\ p_{kk'} & \text{if } b = a + \frac{1}{2} \end{cases}.$$

*Proof.* Straightforward computation verifies the claim. □

Next, we generalize the graded quiver varieties of [Nak11] from the bipartite to the acyclic case, modifying the original construction via the weight order defined above.

Consider the opposite quiver  $Q^{op}$  and denote its set of arrows by  $\Omega$ . Define the function  $\epsilon : H \rightarrow \{\pm 1\}$  such that  $\epsilon(\Omega) = \{1\}$ ,  $\epsilon(\bar{\Omega}) = \{-1\}$ . Similarly,

<sup>3</sup>The  $l$  here stands for “loop” in the quantum loop algebras.

for a linear map  $B_h$  indexed by  $h \in H$ , we define  $(\epsilon B)_h = \epsilon(h)B_h$ . Given vectors  $v, v'$  in  $\mathbb{N}^{I \times (\mathbb{Z} + \frac{1}{2})}$ , and  $w$  in  $\mathbb{N}^{I \times \mathbb{Z}}$ , we define the following graded vector spaces

$$\begin{aligned} \mathbf{L}^\bullet(v, v') &= \oplus_{(i,a)} \text{Hom}(V_i(a), V'_i(a)), \\ \mathbf{L}^{<w}(w, v) &= \oplus_{(i,a)} \text{Hom}(W_i(a), V_i(a - \frac{1}{2})), \\ \mathbf{L}^{<w}(v, w) &= \oplus_{(i,a)} \text{Hom}(V_i(a), W_i(a - \frac{1}{2})), \\ \mathbf{E}^{<w}(v, v') &= (\oplus_{h \in \Omega, a} \text{Hom}(V_{s(h)}(a), V'_{t(h)}(a))) \oplus (\oplus_{\bar{h} \in \bar{\Omega}, a} \text{Hom}(V_{s(\bar{h})}(a), V'_{t(\bar{h})}(a - 1))). \end{aligned}$$

Notice that the weight order strictly decreases along the linear maps in the last three spaces.

Consider the affine space

$$(14) \quad \text{Rep}^\bullet(Q^{op}, v, w) = \mathbf{L}^{<w}(w, v) \oplus \mathbf{L}^{<w}(v, w) \oplus \mathbf{E}^{<w}(v, v).$$

Denote the coordinates of its points by

$$(15) \quad \begin{aligned} (B, \alpha, \beta) &= ((B_h)_{h \in H}, \alpha, \beta) = ((b_h)_{h \in \Omega}, (b_{\bar{h}})_{\bar{h} \in \bar{\Omega}}, (\alpha_i)_i, (\beta_i)_i) \\ &= ((\oplus_a b_{h,a})_h, (\oplus_a b_{\bar{h},a})_{\bar{h}}, (\oplus_a \alpha_{i,a})_i, (\oplus_a \beta_{i,a})_i). \end{aligned}$$

Define the analogue of the moment map  $\mu : \text{Rep}^\bullet(Q^{op}, v, w) \rightarrow \mathbf{L}^\bullet(v, v[-1])$  such that

$$(16) \quad \begin{aligned} \mu(B, \alpha, \beta) &= \oplus_{a \in \mathbb{Z} + \frac{1}{2}, i \in I} \mu(B, \alpha, \beta)_{i,a} \\ &= \oplus_{i,a} (\sum_{h \in \Omega} (b_{h,a} b_{\bar{h},a+1} - b_{\bar{h},a+1} b_{h,a+1}) + \alpha_{i,a+\frac{1}{2}} \beta_{i,a+1}), \end{aligned}$$

or  $\mu(B, \alpha, \beta) = (\epsilon B)B + \alpha\beta$  for short.


**Example 4.1.7.** Let  $Q$  be given as in Figure 1. Figure 5 is an example of the vector space  $\text{Rep}^\bullet(Q^{op}, v, w)$ , where the vertices denote the  $(i, a)$ -degree components of  $W$  and  $V$  and the arrows denote the corresponding linear maps. The components in the same rows have the same  $i$ -degrees, and those in the same columns have the same  $a$ -degrees (or degrees for short).

The group  $G_v = \prod_{i,a} GL(V_{i,a})$  acts naturally on the level set  $\mu^{-1}(0)$  such that for  $g = (g_i)_i = (\oplus_a g_{i,a})_i \in G_v$ ,  $g(\alpha) = g\alpha$ ,  $g(\beta) = \beta g^{-1}$ ,  $g(b_h) = g_{t(h)} b_h g_{s(h)}^{-1}$ ,  $g(b_{\bar{h}}) = g_{t(\bar{h})} b_{\bar{h}} g_{s(\bar{h})}^{-1}$ . Let  $\chi$  be the character of  $G_v$  such that  $\chi(g) = \prod_{i,a} (\det g_{i,a})^{-1}$ .

Define  $\mathcal{M}^\bullet(v, w)$  to be the GIT quotient of  $\mu^{-1}(0)$  with respect to  $\chi$  and  $\mathcal{M}_0^\bullet(v, w)$  to be the categorical quotient of  $\mu^{-1}(0)$  by the action of  $G_v$ . There is a natural projective morphism  $\pi$  from  $\mathcal{M}^\bullet(v, w)$  to  $\mathcal{M}_0^\bullet(v, w)$ . Denote the fibre of  $\pi$  over a point  $x \in \mathcal{M}_0^\bullet(v, w)$  by  $\mathbf{m}^\bullet_x(v, w)$ . When  $x = 0$ , we also denote the fibre by  $\mathcal{L}^\bullet = \mathcal{L}^\bullet(v, w)$ . The varieties  $\mathcal{M}^\bullet(v, w)$ ,  $\mathcal{M}_0^\bullet(v, w)$ ,  $\mathcal{L}^\bullet(v, w)$  are called *graded quiver varieties*.

**Remark 4.1.8.** If the quiver  $Q^{op}$  is bipartite, modulo appropriate degree shifts, our  $q$ -Cartan matrix  $C_q$  and graded quiver varieties agree with those defined in [Nak11].

For the rest of this section, we present important properties of graded quiver varieties.

FIGURE 5. Vector space  $\text{Rep}^\bullet(Q^{op}, v, w)$ 

**4.2. Ungraded quiver varieties.** Let  $V$  and  $W$  be finite-dimensional  $I$ -graded complex vector spaces (without  $a$ -grading). In analogy with the previous subsection, we have vector spaces

$$\begin{aligned} L(V, V) &= \oplus_i \text{Hom}(V_i, V_i) \\ L(W, V) &= \oplus_i \text{Hom}(W_i, V_i) \\ L(V, W) &= \oplus_i \text{Hom}(V_i, W_i) \\ E(V, V) &= \oplus_{h \in H} \text{Hom}(V_{s(h)}, V_{t(h)}). \end{aligned}$$

Consider the symplectic vector space  $\text{Rep}(Q^{op}, V, W) = L(W, V) \oplus L(V, W) \oplus E(V, V)$ . The associated moment map  $\mu : \text{Rep}(Q^{op}, V, W) \rightarrow L(V, V)$  takes a point  $(B, \alpha, \beta)$  of  $\text{Rep}(Q^{op}, V, W)$  to

$$\mu(B, \alpha, \beta) = (\epsilon B)B + \alpha\beta.$$

Following the arguments of [Nak98], we consider the  $GL(V)$ -variety  $\mu^{-1}(0)$ , and fix the character  $\chi$  of  $GL(V)$  such that  $\chi(g) = \prod_i (\det g_i)^{-1}$ . Then we can construct the geometric invariant theory quotient (GIT quotient for short)  $\mathcal{M}(V, W)$  with respect to  $\chi$  and the categorical quotient  $\mathcal{M}_0(V, W)$  by the action of  $GL(V)$  together with the projective morphism  $\pi : \mathcal{M}(V, W) \rightarrow \mathcal{M}_0(V, W)$ .

The points in  $\mathcal{M}_0(V, W)$  are in bijection with the closed orbits in  $\mu^{-1}(0)$ . A point  $(B, \alpha, \beta)$  in such a closed orbit is called a representative of the corresponding point in  $\mathcal{M}_0(V, W)$ , which is denoted by  $[B, \alpha, \beta]$ . Let  $\mu^{-1}(0)^s$  be the open subset of  $\mu^{-1}(0)$  consisting of the  $\chi$ -stable points. It is well known that  $GL(V)$  acts freely on  $\mu^{-1}(0)^s$ . Therefore, the points of  $\mathcal{M}(V, W)$  are in bijection with the orbits of  $\mu^{-1}(0)^s$ . Again, a point  $(B, \alpha, \beta)$  of such a free orbit is called a representative of the corresponding point in  $\mathcal{M}(V, W)$ , denoted by  $[B, \alpha, \beta]$ .

**Proposition 4.2.1** ([Nak98, Corollary 3.12]). *The variety  $\mathcal{M}(V, W)$  is smooth.*

Given any two vectors  $v, v'$  such that  $v \geq v'$  with respect to the dominance order, there is a natural embedding of  $\mathcal{M}_0(V, W)$  into  $\mathcal{M}_0(V', W)$  given by extending the coordinates of the representatives by zero. Take all possible  $v$  and define  $\mathcal{M}_0(W) = \cup_V \mathcal{M}_0(V, W)$  to be the direct limit of all the embeddings. It is possibly infinite dimensional, cf. [Nak01a, 2.5].

Let  $[B, \alpha, \beta]$  be a point in  $\mathcal{M}(V, W)$  and let  $x = (B, \alpha, \beta)$  be its representative. Suppose that we have a  $B$ -invariant filtration of  $V$

$$0 \subset F^0 \subset F^1 \subset \dots \subset F^t = V,$$

where  $\text{Im } \alpha \subset F^0$ . Let  $\text{gr}_0 \alpha$  denote the morphism from  $W$  to  $F^0$  such that its composition with the inclusion  $F^0 \rightarrow V$  is  $\alpha$ . Let  $\text{gr}_0 \beta$  denote the restriction of  $\beta$  to  $F^0$ . For  $1 \leq s \leq t$ , let  $\text{gr}_s B$  denote the endomorphism which  $B$  induces on  $F^s/F^{s-1}$  and  $\text{gr} B = \oplus_{1 \leq s \leq t} \text{gr}_s B$  the endomorphism on  $\oplus_{1 \leq s \leq t} F^s/F^{s-1}$ . The induced representative  $\text{gr} x$  is defined to be  $(\text{gr} B, \text{gr}_0 \alpha, \text{gr}_0 \beta)$ , cf. [Nak98, Definition 3.19].

**Proposition 4.2.2.** [Nak98, Proposition 3.20] *Let  $[B, \alpha, \beta]$  be a point in  $\mathcal{M}(V, W)$  and let  $x = (B, \alpha, \beta)$  be its representative. Then there exists a  $B$ -invariant filtration of  $V$*

$$0 \subset F^0 \subset F^1 \subset \dots \subset F^t = V,$$

*such that  $\text{Im } \alpha \subset F^0$  and the induced triple  $\text{gr} x = (\text{gr} B, \text{gr}_0 \alpha, \text{gr}_0 \beta)$  is a representative of  $\pi([B, \alpha, \beta])$ .*

If  $\hat{G}$  is a subgroup of  $G_v$ , we denote by  $(\hat{G})$  the conjugacy class of  $\hat{G}$ . There is a natural stratification  $\mathcal{M}_0(V, W) = \sqcup_{(\hat{G})} \mathcal{M}_0(V, W)_{(\hat{G})}$ , such that each stratum is the set of the points  $[B, \alpha, \beta]$  which have representatives  $(B, \alpha, \beta)$  with the stabilizers in the conjugacy class  $(\hat{G})$ .

**Proposition 4.2.3** (3.27, [Nak98]). *Let  $[B, \alpha, \beta]$  be a point in  $\mathcal{M}_0(V, W)_{(\hat{G})}$  for some nontrivial  $\hat{G}$ . Then there is a representative  $(B, \alpha, \beta)$  and a decomposition*

$$V = V^0 \oplus (V^1)^{\oplus \hat{v}_1} \oplus \dots \oplus (V^r)^{\oplus \hat{v}_r},$$

*such that we have*

- (1)  $B(V^s) \subset V^s$  for each summand  $V^s$ ,  $0 \leq s \leq r$ ;
- (2) if  $s \neq s'$ , there is no isomorphism from  $V^s$  to  $V^{s'}$  that commutes with  $B$ ;
- (3)  $\text{Im } \alpha$  is contained in  $V^0$ , and  $V^s$  is contained in  $\text{Ker } \beta$  for all  $s > 0$ ;
- (4) the restriction of  $(B, \alpha, \beta)$  to  $V^0$  has the trivial stabilizer in  $\prod_{i \in I} GL(V_i^0)$ ;
- (5) the subgroup  $\prod_{i \in I} GL(V_i^s)$  meets  $\hat{G}$  only in the scalar subgroup  $\mathbb{C}^* \subset GL(V)$ ;

**Remark 4.2.4.** *Restricting the equation  $\mu(B, \alpha, \beta) = 0$  to each summand  $V^s$ ,  $s > 0$ , we see that  $V^s$  is a module over the preprojective algebra associated with  $Q^{\text{op}}$  with respect to the restriction of the  $B$ -action. It has the minimal possible stabilizer  $\mathbb{C}^*$  and a closed orbit under the  $\prod_i GL(V_i^s)$  action. Therefore, it is a representative of a point in the categorical quotient  $\mathcal{M}_0(V^s, 0)$ .*

We call  $\mathcal{M}_0(V, W)_{(\{1\})}$  the regular stratum and denote it by  $\mathcal{M}_0^{\text{reg}}(V, W)$ . It is known that the restriction of  $\pi$  gives an isomorphism from  $\pi^{-1}(\mathcal{M}_0^{\text{reg}}(V, W))$  to  $\mathcal{M}_0^{\text{reg}}(V, W)$ .

Assume  $x$  is a point in  $\mathcal{M}_0^{\text{reg}}(V^0, W)$ , which is naturally embedded into a quotient  $\mathcal{M}_0(V, W)$ . Let  $T$  be the tangent space of  $\mathcal{M}_0^{\text{reg}}(V, W)$  at  $x$ . Since  $\mathcal{M}_0^{\text{reg}}(V^0, W)$  is non-empty,  $(V^0, W)$  is  $l$ -dominant. Define  $W^\perp = \mathbb{C}^{\dim W - \dim V^0}$ ,  $V^\perp = \mathbb{C}^{\dim V - \dim V^0}$ . We have the following theorem

**Theorem 4.2.5** (Theorem 3.3.2, [Nak04]). *There exist neighborhoods  $U$ ,  $U_T$ ,  $U^\perp$  of  $x \in \mathcal{M}_0(V, W)$ ,  $0 \in T$ ,  $0 \in \mathcal{M}_0(V^\perp, W^\perp)$  respectively, and biholomorphic maps  $U \rightarrow U_T \times U^\perp$ ,  $\pi^{-1}(U) \rightarrow U_T \times \pi^{-1}(U^\perp)$ , such that the following diagram commutes:*

$$\begin{array}{ccc} \mathcal{M}(V, W) \supset \pi^{-1}(U) & \xrightarrow[\cong]{} & U_T \times \pi^{-1}(U^\perp) \subset T \times \mathcal{M}(V^\perp, W^\perp) \\ \pi \downarrow & & \downarrow \mathbf{1} \times \pi \\ \mathcal{M}_0(V, W) \supset U & \xrightarrow[\cong]{} & U_T \times U^\perp \subset T \times \mathcal{M}_0(V^\perp, W^\perp) \end{array}$$

In particular, the fibre  $\mathfrak{m}_x(V, W) = \pi^{-1}(x)$  is biholomorphic to the zero fibre  $\mathcal{L}(V^\perp, W^\perp)$  over  $0 \in \mathcal{M}_0(V^\perp, W^\perp)$ .

For any two  $I$ -graded vector spaces  $V, V'$ , define  $L(V, V') = \oplus_i \text{Hom}(V_i, V'_i)$ ,  $E(V, V') = \oplus_h \text{Hom}(V_{s(h)}, V'_{t(h)})$ .

For a point  $(B, \alpha, \beta) \in \mu^{-1}(0)$ , we consider the following complex

$$(17) \quad L(V, V) \xrightarrow{\iota} E(V, V) \oplus L(W, V) \oplus L(V, W) \xrightarrow{d\mu} L(V, V),$$

where  $d\mu$  is the differential map of  $\mu$ , and for  $\xi \in L(V, V)$ , we have

$$(18) \quad \iota(\xi) = (\oplus_{h \in \Omega} (B_h \xi - \xi B_h)) \oplus (-\xi \alpha) \oplus (\beta \xi).$$

Then the tangent space of  $\mathcal{M}(V, W)$  at  $[B, \alpha, \beta]$  is isomorphic to the middle cohomology of this complex.

**4.3. Graded quiver varieties as fixed point sets.** Let  $V$  and  $W$  be as in Section 4.2. Using the fixed point technique developed in [Nak01a], we can deduce the properties of our graded quiver varieties from their ungraded version in Section 4.2.

Choose a torus action<sup>4</sup> of  $\mathbb{C}^*$  on  $\text{Rep}(Q^{op}, V, W)$ , such that for any  $\varepsilon \in \mathbb{C}^*$ , we have

$$\varepsilon(\alpha, \beta, b_h, b_{\bar{h}}) = (\varepsilon^n \alpha, \varepsilon^n \beta, \varepsilon^{2(s(h)-t(h))} b_h, \varepsilon^{2(n-s(h)+t(h))} b_{\bar{h}}).$$

Taking into account of the natural action of  $GL(W)$  on  $\text{Rep}(Q^{op}, V, W)$ , we get a group action of  $GL(W) \times \mathbb{C}^*$  on this space.

Since the actions of  $GL(W) \times \mathbb{C}^*$  and  $GL(V)$  commute,  $GL(W) \times \mathbb{C}^*$  acts on the quiver varieties  $\mathcal{M}(V, W)$  and  $\mathcal{M}_0(V, W)$  and it commutes with the projective morphism  $\pi$ .

Take a pair  $(s, \varepsilon)$ , such that  $s \in GL(W)$  is semisimple, and  $\varepsilon$  is not a root of unity. It generates a cyclic subgroup, whose closure with respect to the Zariski topology is denoted by  $A$ . As in [Nak01a], let  $[B, \alpha, \beta]$  be a

<sup>4</sup>The choice is not unique. In fact, different choices might give isomorphic graded quiver varieties.

point  $\mathcal{M}(V, W)^A$  and  $(B, \alpha, \beta)$  be any representative of it. There exists a group homomorphism  $\rho_{(B, \alpha, \beta)}$  from  $A$  to  $GL(V)$  such that for any element  $a \in A$ , we have  $a(B, \alpha, \beta) = \rho_{(B, \alpha, \beta)}(a)^{-1}(B, \alpha, \beta)$ . The conjugacy class of  $\rho_{(B, \alpha, \beta)}$  is independent of the choice of the representative  $(B, \alpha, \beta)$  because the actions of  $GL(V)$  and  $A$  commute. So we can denote it by  $[\rho_{[B, \alpha, \beta]}]$ .

**Lemma 4.3.1** ([Nak01a, Section 4.1]). *The map from  $\mathcal{M}(V, W)^A$  to the conjugacy classes of the group homomorphisms from  $A$  to  $GL(V)$ , sending  $[B, \alpha, \beta]$  to  $[\rho_{[B, \alpha, \beta]}]$ , is locally constant.*

*Proof.* Since  $A$  is generated by the element  $a = (s, \varepsilon)$ , it suffices to study the conjugacy class  $[\rho_{[B, \alpha, \beta]}(a)]$ .

First, we show that  $[\rho_{[B, \alpha, \beta]}(a)]$  is continuous in  $[B, \alpha, \beta]$ . Recall that  $\mu^{-1}(0)^s$  is a principal  $GL(V)$ -bundle over  $\mathcal{M}(V, W)$ . Denote the fibre map by  $p$ . Take a trivialization. Let  $U$  be any chart. For any continuous curve  $[x(t)]$ ,  $0 \leq t \leq 1$ , the curve  $[x^U(t)] = U \cap [x(t)]$  in  $U \cap \mathcal{M}(V, W)^A$  can be lifted to a continuous curve  $x^U(t) = [x^U(t)] \times \{e\}$  in  $(U \times GL(V)) \cap p^{-1}(\mathcal{M}(V, W)^A)$ , where  $e$  denotes the identity of  $GL(V)$ . Then the fibre coordinates of the continuous curve  $a^{-1}x^U(t)$  are described by  $\rho_{x(t)}(a)$ . Recall that the transition between different charts is given by conjugating. Therefore, the conjugacy class  $[\rho_{[x(t)]}(a)]$  is continuous on the curve  $[x(t)]$ .

Since  $s$  is semi-simple and  $\rho : A \rightarrow GL(V)$  is a group homomorphism, the Jordan form of  $\rho_{x(t)}(a)$  is a discrete subset in the set of the conjugacy classes of  $GL(V)$ . Therefore,  $[\rho_{[x(t)]}(a)]$  is locally constant.  $\square$

We denote the collection of the points  $[B, \alpha, \beta]$  with the common conjugacy class  $[\rho]$  by  $\mathcal{M}([\rho])$ . Then  $\mathcal{M}([\rho])$  is a union of connected components of  $\mathcal{M}(V, W)^A$ . It follows that we have  $\pi \mathcal{M}(V, W)^A = \cup_{[\rho]} \pi \mathcal{M}([\rho]) = \sqcup_{[\rho]} \pi \mathcal{M}([\rho])$ . Denote each stratum  $\pi \mathcal{M}([\rho])$  by  $\mathcal{M}_0([\rho])$ .

Fix the conjugacy class  $[\rho]$ . Using the eigenvalues and the eigenspaces of  $s$  and  $\rho(s, \varepsilon)$ , we can endow  $W$  and  $V$  with gradings. Assume  $W$  and  $V$  have eigenspace decompositions  $W = \oplus_i W_i = \oplus_{i; a \in \mathbb{Z}} W_i(a)$ ,  $V = \oplus_i V_i = \oplus_{i; a \in \mathbb{Z}} V_i(a + \frac{1}{2})$ , where  $W_i(a)$  and  $V_i(a + \frac{1}{2})$  have eigenvalues  $\varepsilon^{2(an+i-1)}$ ,  $\varepsilon^{2((a+\frac{1}{2})n+i-1)}$  respectively. Associate to  $[\rho(s, \varepsilon)]$  the bigraded vectors  $w = (\dim W_i(a))$ ,  $v = (\dim V_i(a))$ . We can identify  $\mathcal{M}([\rho])$  with the graded quiver variety  $\mathcal{M}^\bullet(v, w)$ . Similarly, the graded categorical quotient  $\mathcal{M}_0^\bullet(v, w)$  is identified with the subvariety  $\mathcal{M}_0([\rho])$  of  $\mathcal{M}_0(V, W)$ .

**Remark 4.3.2.** *If we take the  $\mathbb{C}^*$ -action in [Nak11], the representatives of the fixed points in the sub-varieties  $\mathcal{M}([\rho])$  do not take the form of representations of the quiver in Figure 5. For example, let the quiver  $Q$  be given by Figure 1. For simplicity, let us assume  $w_2 = 0$ . Then these representatives are representations of the quiver in Figure 6, where the black arrows arise from those of  $Q^{op}$ , the green arrows arise from those of  $Q$ , and the orange arrows correspond to the linear maps  $\alpha_i(a)$ ,  $\beta_i(a)$ ,  $i \in I$ ,  $a \in \mathbb{Z}$ .*

*Such representations do not suit our purpose.*

The graded version of Proposition 4.2.1 implies that the graded quiver variety  $\mathcal{M}^\bullet(V, W)$  is smooth.




FIGURE 6. The vector space  $\text{Rep}^\bullet(Q^{op}, v, w)$  for  $Q$  acyclic and the  $\mathbb{C}^*$ -action of [Nak11]

**Proposition 4.3.3** ([Nak94, Corollary 5.5] [Nak01a, Proposition 4.1.2]).  $\mathcal{M}([\rho])$  is homotopic to  $\mathcal{L}(V, W) \cap \mathcal{M}([\rho])$ .

*Proof.* By using Slodowy's technique [Slo80, Section 4.3], Nakajima has shown that  $\mathcal{M}(V, W)$  is homotopic to  $\mathcal{L}(V, W)$  in [Nak94], and he has also proved this proposition with a different  $GL(W) \times \mathbb{C}^*$ -action in [Nak01a]. The technique remains valid here. We shall briefly recall it.

Equip the space  $\text{rep}(Q^{op}, V, W)$  with the  $\mathbb{C}^*$ -action such that we have  $\varepsilon(B_h, B_{\bar{h}}, \alpha, \beta) = (B_h, \varepsilon B_{\bar{h}}, \alpha, \varepsilon \beta)$  for any  $\varepsilon \in \mathbb{C}^*$ . It commutes with the  $GL(V)$ -action. Furthermore, the set of the stable points  $\mu^{-1}(0)^s$  is invariant under this action. Therefore, we obtain a  $\mathbb{C}^*$ -action on  $\mathcal{M}(V, W)$ . We have

$$(19) \quad \mathcal{L}(V, W) = \{[x] \in \mathcal{M}(V, W) \mid \lim_{\varepsilon \rightarrow \infty} \varepsilon[x] \text{ exists}\}.$$

Now the technique of Slodowy [Slo80, 4.3] implies that, via this  $\mathbb{C}^*$ -action,  $\mathcal{M}(V, W)$  retracts to a neighborhood of  $\mathcal{L}(V, W)$ , such that  $\mathcal{L}(V, W)$  is a strong deformation retract of this neighborhood.

Because our  $A$ -action commutes with this  $\mathbb{C}^*$ -action, we can apply the above constructions to the  $A$ -fixed subsets  $\text{Rep}^\bullet(Q^{op}, v, w)$ ,  $\mathcal{M}^\bullet(v, w)$ ,  $\mathcal{L}^\bullet(v, w)$ . Then the proposition is verified.  $\square$

Let us define  $\mathcal{M}_0^{\bullet \text{reg}}(v, w) = \mathcal{M}_0^{\bullet \text{reg}}([\rho]) = \pi(\pi^{-1}(\mathcal{M}_0^{\text{reg}}(V, W)) \cap \mathcal{M}([\rho]))$ . Then the morphism  $\pi$  is an isomorphism from  $\pi^{-1}(\mathcal{M}_0^{\bullet \text{reg}}(v, w))$  to  $\mathcal{M}_0^{\bullet \text{reg}}(v, w)$ .

The maps in Theorem 4.2.5 commute with the  $GL(W) \times \mathbb{C}^*$  action. Restrict the maps to the subvarieties  $\mathcal{M}^\bullet(v, w)$ ,  $\mathcal{M}^\bullet(v^\perp, w^\perp)$  of  $\mathcal{M}(V, W)$ ,  $\mathcal{M}(V^\perp, W^\perp)$  respectively. We obtain a transversal slice theorem for graded quiver varieties.

**Theorem 4.3.4** (Transversal slice). *Assume  $x$  is a point in  $\mathcal{M}_0^{\bullet \text{reg}}(v^0, w)$ , which is naturally embedded into a quotient  $\mathcal{M}_0^\bullet(v, w)$ . Let  $T$  be the tangent space of  $\mathcal{M}_0^{\bullet \text{reg}}(v, w)$  at  $x$ . As  $(v^0, w)$  is  $l$ -dominant, define  $w^\perp = w - C_q v^0$ ,  $v^\perp = v - v^0$ . Then there exist neighborhoods  $U$ ,  $U_T$ ,  $U^\perp$  of  $x \in \mathcal{M}_0^\bullet(V, W)$ ,*

$0 \in T$ ,  $0 \in \mathcal{M}_0^\bullet(v^\perp, w^\perp)$  respectively, and biholomorphic maps  $U \rightarrow U_T \times U^\perp$ ,  $\pi^{-1}(U) \rightarrow U_T \times \pi^{-1}(U^\perp)$ , such that the following diagram commutes:

$$\begin{array}{ccc} \mathcal{M}^\bullet(v, w) \supset \pi^{-1}(U) & \xrightarrow{\cong} & U_T \times \pi^{-1}(U^\perp) \subset T \times \mathcal{M}(v^\perp, w^\perp) \\ \pi \downarrow & & \downarrow \mathbf{1} \times \pi \\ \mathcal{M}_0^\bullet(v, w) \supset U & \xrightarrow{\cong} & U_T \times U^\perp \subset T \times \mathcal{M}_0^\bullet(v^\perp, w^\perp) \end{array}$$

In particular, the fibre  $\mathbf{m}_x^\bullet(v, w) = \pi^{-1}(x)$  is biholomorphic to the zero fibre  $\mathcal{L}^\bullet(v^\perp, w^\perp)$  over  $0 \in \mathcal{M}_0^\bullet(v^\perp, w^\perp)$ .

**Proposition 4.3.5.** *We have a stratification*

$$(20) \quad \mathcal{M}_0^\bullet(v, w) = \sqcup_{(v', w) \geq (v, w)} \mathcal{M}_0^{\bullet \text{reg}}(v', w).$$

In particular, the variety  $\mathcal{M}_0^\bullet(w) = \cup_v \mathcal{M}_0^\bullet(v, w)$  has a stratification

$$\mathcal{M}_0^\bullet(w) = \sqcup_v \mathcal{M}_0^{\bullet \text{reg}}(v, w).$$

*Proof.* It suffices to show

$$(21) \quad \pi(\mathcal{M}(V, W)^A) = \sqcup_{V'} \pi(\pi^{-1}(\mathcal{M}_0^{\text{reg}}(V', W)) \cap \mathcal{M}(V, W)^A)$$

for each pair of vector spaces  $(V, W)$ . Then we obtain (20) by restricting (21) to the subvariety  $\mathcal{M}_0^\bullet(v, w) = \mathcal{M}_0([\rho])$  for the conjugacy class  $[\rho]$  associated with  $(v, w)$ .

We claim that every point  $[B, \alpha, \beta]$  of  $\pi(\mathcal{M}(V, W)^A)$  is contained in the right hand side of (21). Using Proposition 4.2.3 and Remark 4.2.4, we see that  $[B, \alpha, \beta]$  belongs to some  $\mathcal{M}_0(V, W)_{(\widehat{G})}$ . If  $\widehat{G}$  equals  $\{e\}$ , the claim is true. Otherwise, choose the representative  $(B, \alpha, \beta)$  in Proposition 4.2.3 and consider the  $B$  actions on all the summand  $V^s$ ,  $s > 0$ . If the actions are trivial,  $[B, \alpha, \beta]$  lies in the regular stratum  $\mathcal{M}_0^{\text{reg}}(V^0, W)$ , and the claim follows easily. If the action is nontrivial for some  $V^s$ ,  $s > 0$ , we obtain a point other than 0 in the categorical quotient  $\mathcal{M}_0(V^s, 0)$ . Because the  $GL(W) \times \mathbb{C}^*$ -action is compatible with the decomposition of  $V$  in Proposition 4.2.3,  $\rho_{(B, \alpha, \beta)}$  stabilizes the decomposition. Let  $v^s$  be the bigraded vector associated with the  $\rho_{(B, \alpha, \beta)}$ -action on  $V^s$ . As in Remark 4.2.4,  $V^s$  is a representative of a nonzero point in  $\mathcal{M}_0^\bullet(v^s, 0)$ . However in our setting  $\mathcal{M}_0^\bullet(v^s, 0)$  is the categorical quotient of the  $v^s$ -dimensional representations of some acyclic quiver, which is always equal to  $\{0\}$ . This contradiction implies that the  $B$ -action on  $V^s$  must be trivial.  $\square$

**Remark 4.3.6.** *When  $Q$  is of Dynkin type, the ungraded version of the proposition holds, cf. [Nak01a, Proposition 2.6.3]. However, it does not necessarily hold when  $Q$  is not of Dynkin type, cf. Example 10.10 in [Nak98]. In general, whether the proposition is true or not depends on the choice of the  $GL(W) \times \mathbb{C}^*$ -action.*

Let  $m$  be any integer. Following [Nak01a, Section 2.8], we define a  $\mathbb{C}^*$ -module structure  $L(m)$  on  $\mathbb{C}$  by

$$(22) \quad \varepsilon \cdot v = \varepsilon^m v,$$

where  $\varepsilon \in \mathbb{C}^*$ ,  $v \in \mathbb{C}$ .

For a  $\mathbb{C}^*$ -module  $M$ , we denote the  $\mathbb{C}^*$ -module  $L(m) \otimes_{\mathbb{C}} M$  by  $q^m M$ .

As in [2.9, [Nak01a]],  $\mu^{-1}(0)^s$  is a principal  $GL(V)$ -bundle over  $\mathcal{M}(V, W)$ . Therefore, for any  $i \in I$ , we can view the vector space  $V_i$  as an associated vector bundle by using the natural  $GL(V_i)$  action. Also, it is naturally a  $GL(W) \times \mathbb{C}^*$ -equivariant vector bundle such that  $GL(W)$  acts trivially. Similarly, we view  $W_i$  as a  $GL(W) \times \mathbb{C}^*$ -equivariant vector bundle over  $\mathcal{M}(V, W)$  by using the  $GL(W_i) \times \mathbb{C}^*$  action such that  $\mathbb{C}^*$  acts trivially.

For each  $k \in I$ , we have the following complex  $C_k^* = (\sigma_k, \tau_k)$  of  $GL(W) \times \mathbb{C}^*$ -equivariant vector bundles:

(23)

$$C_k^* : q^{-2n}V_k \xrightarrow{\sigma_k} ((\oplus_{j:i < k} q^{2(k-i-n)} V_i^{\oplus b_{ik}}) \oplus (\oplus_{j:j > k} q^{2(j-k)} V_j^{\oplus b_{kj}}) \oplus q^{-n}W_k) \xrightarrow{\tau_k} V_k,$$

where  $\sigma_k = (\oplus_{h \in H, s(h)=k} B_h) \oplus \beta_k$ ,  $\tau_k = \sum_{h \in \Omega, t(h)=k} B_h - \sum_{\bar{h} \in \bar{\Omega}, t(\bar{h})=k} B_{\bar{h}} + \alpha_k$  are  $GL(W) \times \mathbb{C}^*$ -equivariant morphisms. This complex is just the complex in [Nak98][4.2] with a modification of  $GL(W) \times \mathbb{C}^*$ -action. Let the middle term be the degree 0 component.

**Proposition 4.3.7** ([Nak01a, Lemma 2.9.2, Lemma 2.9.4]). *Fix a point  $[B, \alpha, \beta] \in \mathcal{M}(V, W)$  and consider  $C_k^*$  as a complex of vector spaces,  $k \in I$ .*

- 1) *The cohomology  $H^{-1}(C_k^*)$  vanishes.*
- 2) *If the image  $\pi([B, \alpha, \beta]) \in \mathcal{M}_0(V, W)$  is contained in some regular stratum  $\mathcal{M}_0^{\text{reg}}(V', W) \subset \mathcal{M}_0(V, W)$ , then  $V'$  equals  $V$  if and only if  $H^1(C_k^*)$  vanishes for all  $k \in I$ .*

Fix a graded quiver variety  $\mathcal{M}^\bullet(v, w) = \mathcal{M}([\rho])$ . Let  $(C_k^*)^\bullet$  be the restriction of  $C_k^*$  to  $\mathcal{M}^\bullet(v, w)$ . Then the complex  $(C_k^*)^\bullet$  decomposes as

$$(24) \quad (C_k^*)^\bullet = \oplus_{b \in \mathbb{Z} + \frac{1}{2}} C_k^*(b),$$

where the complexes  $C_k^*(b) = (\sigma_k(b), \tau_k(b))$  are given by

$$(25) \quad q^{-2n}V_k(b) \xrightarrow{\sigma_k(b)} ((\oplus_{j:i < k} q^{2(k-i-n)} V_i(b)^{\oplus b_{ik}}) \oplus (\oplus_{j:j > k} q^{2(j-k)} V_j(b-1)^{\oplus b_{kj}}) \oplus q^{-n}W_k(b - \frac{1}{2})) \xrightarrow{\tau_k(b)} V_k(b-1).$$

The decomposition commutes with the  $A$ -action.

We have the following analogue of Proposition 4.3.7 in graded cases.

**Proposition 4.3.8.** 1) *The cohomology  $H^{-1}((C_k^*)^\bullet)$  vanishes.*

- 2) *The image  $\pi([B, \alpha, \beta]) \in \mathcal{M}_0^\bullet(v, w)$  is contained in the regular stratum  $\mathcal{M}_0^{\bullet, \text{reg}}(v, w)$  if and only if  $H^1((C_k^*)^\bullet)$  vanishes,  $\forall k \in I$ .*

*Proof.* Part 1) follows from Proposition 4.3.7 by restriction. For part 2), we additionally use Proposition 4.3.5.  $\square$

**Theorem 4.3.9.**  $\mathcal{M}^\bullet(v, w)$  is connected.

*Proof.* The statement follows from the arguments in the proof of [Nak01a, Theorem 5.5.6]. Notice that, since our  $GL(W) \times \mathbb{C}^*$ -action is different from that of [Nak01a], we don't need the condition  $|b_{ij}| \leq 1$ ,  $1 \leq i, j \leq n$ , in [Nak01a, Theorem 5.5.6].  $\square$

As a consequence, the smooth variety  $\mathcal{M}^\bullet(v, w)$  is irreducible.

**Proposition 4.3.10** ([Nak01a, Corollary 5.5.5]). *On a nonempty open subset of  $\mathcal{M}^\bullet(v, w)$ , we have*

$$(26) \quad \text{codim } \text{Im } \tau_k(b) = \max(0, -\text{rank}(C_k^*(b))).$$

**Lemma 4.3.11.**  *$(\text{rank}(C_k^*))_{k \in I}$  equals  $w - C_q v$ .*

**Proposition 4.3.12.**  *$\mathcal{M}_0^{\bullet \text{reg}}(v, w)$  is non-empty if and only if  $(v, w)$  is  $l$ -dominant.*

*Proof.* This follows from Proposition 4.3.8 4.3.10 and Lemma 4.3.11.  $\square$

**Remark 4.3.13.** *For any given vector  $w$ , by induction on its width with respect to the weight order, one can prove that there are only finitely many  $v$  such that  $(v, w)$  is  $l$ -dominant. Then Propositions 4.3.12 and 4.3.5 imply that  $\mathcal{M}_0^\bullet(w) = \cup_v \mathcal{M}_0^\bullet(v, w)$  is finite dimensional.*

**Corollary 4.3.14.** *Let  $(v, w), (v^0, w)$  be  $l$ -dominant pairs. Then we have  $\mathcal{M}_0^{\bullet \text{reg}}(v^0, w) \subset \overline{\mathcal{M}_0^{\bullet \text{reg}}(v, w)}$  if and only if  $(v^0, w) \geq (v, w)$ .*

*Proof.* Proposition 4.3.12 implies that both regular strata are non-empty. By Theorem 4.3.9,  $\mathcal{M}_0^\bullet(v, w)$  is irreducible. Since  $\mathcal{M}_0^{\bullet \text{reg}}(v, w)$  is a non-empty open subset in  $\mathcal{M}_0^\bullet(v, w)$ , its closure equals  $\mathcal{M}_0^\bullet(v, w)$ .

If  $(v^0, w) \geq (v, w)$ ,  $\mathcal{M}_0^\bullet(v^0, w)$  is naturally embedded into  $\mathcal{M}_0^\bullet(v, w)$ . Thus the only if part holds.

Conversely, if  $\mathcal{M}_0^{\bullet \text{reg}}(v^0, w) \subset \overline{\mathcal{M}_0^{\bullet \text{reg}}(v, w)} = \mathcal{M}_0^\bullet(v, w)$ , using proposition 4.3.5 we obtain the if part.  $\square$

Following [Nak04, (4.9)], for any given pairs of vectors  $(v^1, w^1)$  and  $(v^2, w^2)$ , we define the following complex of vector bundles over  $\mathcal{M}^\bullet(v^1, w^1) \times \mathcal{M}^\bullet(v^2, w^2)$ :

$$(27) \quad \mathbf{L}^\bullet(v^1, v^2) \xrightarrow{\sigma^{21}} \mathbf{E}^{<w}(v^1, v^2) \oplus \mathbf{L}^{<w}(w^1, v^2) \oplus \mathbf{L}^{<w}(v^1, w^2) \xrightarrow{\tau^{21}} \mathbf{L}^\bullet(v^1, v^2[-1]),$$

where the middle term has degree 0, and we denote

$$(28) \quad \sigma^{21}(\xi) = (B^2\xi - \xi B^1) \oplus (-\xi\alpha^1) \oplus \beta^2\xi,$$

$$(29) \quad \tau^{21}(C \oplus I \oplus J) = \epsilon B^2 C + \epsilon C B^1 + \alpha^2 J + I \beta^1.$$

The complex is exact on the left and on the right, cf. the argument in [Nak98, 3.10].

**Lemma 4.3.15.** *The quotient  $\ker \tau^{21} / \text{Im } \sigma^{21}$  is a vector bundle over  $\mathcal{M}^\bullet(V^1, W^1) \times \mathcal{M}^\bullet(V^2, W^2)$ .*

*Proof.* By [CBJ05], any subvariety of a vector bundle over the smooth variety  $\mathcal{M}^\bullet(v^1, w^1) \times \mathcal{M}^\bullet(v^2, w^2)$ , whose intersection with every fibre is a vector subspace of constant dimension, is a sub-bundle. Therefore,  $\ker \tau^{21}$  is a vector bundle. Let  $(\sigma^{21})^*$  be the transpose (or dual) of  $\sigma^{21}$  restricted to  $\ker \tau^{21}$ . Its kernel  $(\ker \tau^{21} / \text{Im } \sigma^{21})^*$  is a sub-bundle of  $(\ker \tau^{21})^*$ . Therefore,  $\ker \tau^{21} / \text{Im } \sigma^{21}$  is again a vector bundle.  $\square$

Denote the rank of the vector bundle  $\ker \tau^{21} / \text{Im } \sigma^{21}$  by  $d((v^1, w^1), (v^2, w^2))$ .

**Lemma 4.3.16.** *The rank  $d((v^1, w^1), (v^2, w^2))$  is given by*

$$(30) \quad d((v^1, w^1), (v^2, w^2)) = (w^1 - C_q v^1) \cdot v^2[-\frac{1}{2}] + v^1 \cdot w^2[-\frac{1}{2}].$$

*Proof.* It suffices to calculate the rank of the complex. We have

$$\begin{aligned}
d((v^1, w^1), (v^2, w^2)) &= \sum_{k \in I, a \in \mathbb{Z}} v_k^2(a - \frac{1}{2}) \left( \sum_{i: i < k} b_{ik} v_i^1(a + \frac{1}{2}) + \sum_{j: j > k} b_{kj} v_j^1(a - \frac{1}{2}) \right) \\
&\quad + \sum_{k \in I, a \in \mathbb{Z}} (w_k^1(a) v_k^2(a - \frac{1}{2}) + v_k^1(a - \frac{1}{2}) w_k^2(a - 1)) \\
&\quad - \sum_{k \in I, a \in \mathbb{Z}} v_k^2(a - \frac{1}{2}) (v_k^1(a - \frac{1}{2}) + v_k^1(a + \frac{1}{2})) \\
&= \sum_{k, a} (v_k^2(a - \frac{1}{2}) (-C_q v^1)_k(a) + w_k^1(a) v_k^2(a - \frac{1}{2}) + w_k^2(a) v_k^1(a + \frac{1}{2})) \\
&= -v^2[-\frac{1}{2}] \cdot C_q v^1 + w^1 \cdot v^2[-\frac{1}{2}] + w^2 \cdot v^1[\frac{1}{2}] \\
&= \text{RHS}.
\end{aligned}$$

□

**Remark 4.3.17.** Using equation (12), we can easily rewrite the quadratic form (30) as

$$(31) \quad d((v^1, w^1), (v^2, w^2)) = v^1[\frac{1}{2}] \cdot (w^2 - C_q v^2) + w^1[\frac{1}{2}] \cdot v^2.$$

The reader is invited to compare this expression with [Nak04, (2.1)].

Let  $[B, \alpha, \beta]$  be any point in  $\mathcal{M}^\bullet(v, w) = \mathcal{M}([\rho]) \subset \mathcal{M}(V, W)$ . The tangent space of  $\mathcal{M}^\bullet(v, w)$  at  $[B, \alpha, \beta]$  is the  $A$ -fixed part of the tangent space of  $\mathcal{M}(V, W)$  at the same point. The  $A$ -fixed part of complex (17) over this point is just

$$(32) \quad \mathbb{L}^{<w}(v, v[-1]) \xrightarrow{\iota} \mathbb{E}^{<w}(v, v) \oplus \mathbb{L}^{<w}(v, w) \oplus \mathbb{L}^{<w}(w, v) \xrightarrow{d\mu} \mathbb{L}^{<w}(v, v[1]).$$

Comparing it with the complex (27), we have the following result.

**Lemma 4.3.18.** The dimension of  $\mathcal{M}^\bullet(v, w)$  equals  $d((v, w), (v, w))$ .

**Theorem 4.3.19** ([Nak01a, Theorem 7.4.1]). The odd homology of  $\mathcal{L}^\bullet(v, w)$  vanishes. And we have a nondegenerate pairing between  $H_*(\mathcal{L}^\bullet(v, w))$  and  $H_*(\mathcal{M}^\bullet(v, w))$ .

*Proof.* We refer the reader to [Nak01a, Section 7] for the proof. This theorem is just a consequence of [Nak01a, Theorem 7.4.1]. □

## 5. $qt$ -CHARACTERS

In the study of finite dimensional representations of quantum affine algebras, Nakajima invented the  $t$ -analogues of the  $q$ -characters ( $qt$ -characters for short) [Nak04], which are defined as the generating series of the Betti numbers of the graded quiver varieties. In this section, we generalize his constructions for our graded quiver varieties and introduce a new  $t$ -deformation.

**5.1. Quadratic forms.** For any two pairs of vectors  $(v^1, w^1), (v^2, w^2)$ , recall that we have the quadratic form  $d((v^1, w^1), (v^2, w^2))$  given by (30), whose geometric meaning is explained by Lemma 4.3.18. Let us further introduce quadratic forms  $\tilde{d}'((v^1, w^1), (v^2, w^2))$ ,  $\tilde{d}'_W(w^1, w^2)$ , and  $\mathcal{E}'(w^1, w^2)$  such that

$$(33) \quad \tilde{d}'_W(w^1, w^2) = -w^1 \left[ \frac{1}{2} \right] \cdot C_q^{-1} w^2,$$

$$(34) \quad \mathcal{E}'(w^1, w^2) = -w^1 \left[ \frac{1}{2} \right] \cdot C_q^{-1} w^2 + w^2 \left[ \frac{1}{2} \right] \cdot C_q^{-1} w^1.$$

$$(35) \quad \tilde{d}'((v^1, w^1), (v^2, w^2)) = d((v^1, w^1), (v^2, w^2)) + \tilde{d}'_W(w^1, w^2),$$

Notice that  $\tilde{d}'((0, w^1), (0, w^2))$  equals  $\tilde{d}'_W(w^1, w^2)$ .

**Remark 5.1.1.** Our  $\tilde{d}'$  and  $\tilde{d}'_W$  are different from  $\tilde{d}$  and  $\tilde{d}_W$  in [Nak04], but the properties are similar.

**Lemma 5.1.2.** The following equality holds:

$$(36) \quad \tilde{d}'((v^1, w^1), (v^2, w^2)) = \tilde{d}'((0, w^1 - C_q v^1), (0, w^2 - C_q v^2)).$$

*Proof.* We have

$$\begin{aligned} \tilde{d}'((v^1, w^1), (v^2, w^2)) &= (w^1 - C_q v^1) \cdot v^2 \left[ -\frac{1}{2} \right] + v^1 \cdot w^2 \left[ -\frac{1}{2} \right] - w^1 \left[ \frac{1}{2} \right] \cdot C_q^{-1} w^2 \\ &= (w^1 - C_q v^1) \cdot v^2 \left[ -\frac{1}{2} \right] + v^1 \cdot C_q C_q^{-1} w^2 \left[ -\frac{1}{2} \right] - w^1 \left[ \frac{1}{2} \right] \cdot C_q^{-1} w^2 \\ &\stackrel{(12)}{=} (w^1 - C_q v^1) \cdot v^2 \left[ -\frac{1}{2} \right] + C_q v^1 \cdot C_q^{-1} w^2 \left[ -\frac{1}{2} \right] - w^1 \left[ \frac{1}{2} \right] \cdot C_q^{-1} w^2 \\ &= -(w^1 - C_q v^1) \left[ \frac{1}{2} \right] \cdot C_q^{-1} (w^2 - C_q v^2) \\ &= \tilde{d}'((0, w^1 - C_q v^1), (0, w^2 - C_q v^2)). \end{aligned}$$

□

**Lemma 5.1.3.** 1) For any  $a \in \mathbb{Z}$  and  $i, j \in I$ , we have

$$(37) \quad \mathcal{E}'(e_{i,a}, e_{j,a}) = 0,$$

$$(38) \quad \mathcal{E}'(e_{i,a}, e_{j,a-1}) = -p_{ji},$$

$$(39) \quad \mathcal{E}'(e_{i,a-1} + e_{i,a}, e_{j,a-1} + e_{j,a}) = p_{ij} - p_{ji}.$$

2) Let  $B_{\tilde{Q}}$  be the  $2n \times 2n$ -matrix associated with  $\tilde{Q}_1^z$  the level 1 ice quiver with  $z$ -pattern whose principal part is  $Q$ . Define  $\Lambda^z$  to be the  $2n \times 2n$  matrix such that for  $1 \leq i, j \leq n$ , we have

$$(40) \quad \Lambda^z_{i,j} = -\mathcal{E}'(e_{i,0}, e_{j,0}) = 0,$$

$$(41) \quad \Lambda^z_{i+n,j} = -\mathcal{E}'(e_{i,-1} + e_{i,0}, e_{j,0}) = -p_{ij},$$

$$(42) \quad \Lambda^z_{i,j+n} = -\mathcal{E}'(e_{i,0}, e_{j,-1} + e_{j,0}) = p_{ji},$$

$$(43) \quad \Lambda^z_{i+n,j+n} = -\mathcal{E}'(e_{i,-1} + e_{i,0}, e_{j,-1} + e_{j,0}) = -p_{ij} + p_{ji}.$$

Then  $\Lambda^z$  is the inverse of  $-B_{\tilde{Q}}$ .

*Proof.* 1) The claim follows from (13).

2) Apply the results of 1) for  $a = 0$ . The entries of  $B_{\tilde{Q}}$  are given by

$$\begin{aligned} (B_{\tilde{Q}})_{i,j} &= b_{ij}, \\ (B_{\tilde{Q}})_{i+n,j} &= -\delta_{ij} + [b_{ji}]_+, \\ (B_{\tilde{Q}})_{i,j+n} &= -(B_{\tilde{Q}})_{j+n,i}, \\ (B_{\tilde{Q}})_{i+n,j+n} &= 0, \end{aligned}$$

for any  $1 \leq i, j \leq n$ . Here  $[\ ]_+$  is defined to be  $\max\{\ , 0\}$ .

We have, for any  $1 \leq i, k \leq n$ ,

$$\begin{aligned} (-\Lambda^z B_{\tilde{Q}})_{i,k} &= \sum_{1 \leq j \leq n} \Lambda^z_{i,j+n} \cdot (-B_{\tilde{Q}})_{j+n,k} \\ &= \sum_{1 \leq j \leq n} p_{ji} \delta_{jk} - \sum_{1 \leq j \leq n} p_{ji} [b_{kj}]_+ \\ &= p_{ki} - \sum_j [b_{kj}]_+ p_{ji} \\ &= \delta_{ik}, \end{aligned}$$

$$\begin{aligned} (-\Lambda^z B_{\tilde{Q}})_{i+n,k} &= \sum_{1 \leq j \leq n} (-\Lambda^z_{i+n,j}) \cdot (B_{\tilde{Q}})_{j,k} + \sum_{1 \leq j \leq n} (-\Lambda^z_{i+n,j+n}) \cdot (B_{\tilde{Q}})_{j+n,k} \\ &= \sum_{1 \leq j \leq n} p_{ij} b_{jk} + \sum_{1 \leq j \leq n} (p_{ij} - p_{ji})(-\delta_{jk} + [b_{kj}]_+) \\ &= \sum_{j:j>k} p_{ij}(-b_{kj}) + \sum_{j:j<k} p_{ij} b_{jk} + (-p_{ik} + p_{ki}) \\ &\quad + \sum_j (p_{ij} - p_{ji}) [b_{kj}]_+ \\ &= \left( \sum_{j:j>k} p_{ij}(-b_{kj}) + \sum_j p_{ij} [b_{kj}]_+ \right) + \left( \sum_{j:j<k} p_{ij} b_{jk} - p_{ik} \right) \\ &\quad + (p_{ki} - \sum_j [b_{kj}]_+ p_{ji}) \\ &= 0 - \delta_{ik} + \delta_{ik} \\ &= 0, \end{aligned}$$

$$(-\Lambda^z B_{\tilde{Q}})_{i,k+n} = (\Lambda^z B_{\tilde{Q}})_{k+n,i} = 0,$$

$$\begin{aligned} (-\Lambda^z B_{\tilde{Q}})_{i+n,k+n} &= \sum_{1 \leq j \leq n} p_{ij} (\delta_{kj} - [b_{jk}]_+) \\ &= p_{ik} - \sum_j p_{ij} [b_{jk}]_+ \\ &= \delta_{ik}. \end{aligned}$$

□

**5.2. Deformed Grothendieck ring.** Recall that, by Proposition 4.3.5, we have the stratification

$$\mathcal{M}_0^\bullet(w) = \cup_v \mathcal{M}_0^\bullet(v, w) = \sqcup_{v: (v, w) \text{ is } l\text{-dominant}} \mathcal{M}_0^{\bullet\text{reg}}(v, w).$$

For any given pair  $(v, w)$ , the map  $\pi : \mathcal{M}^\bullet(v, w) \rightarrow \mathcal{M}_0^\bullet(w)$  is proper since it is the composition of a projective morphism and a closed embedding. The rank-1 trivial local system over the smooth variety  $\mathcal{M}^\bullet(v, w)$  yields a perverse sheaf  $1_{\mathcal{M}^\bullet(v, w)} = \mathbb{C}[\dim \mathcal{M}^\bullet(v, w)]$ . If  $(v, w)$  is  $l$ -dominant,  $\mathcal{M}_0^{\bullet\text{reg}}(v, w)$  is nonempty by Proposition 4.3.12. Define  $IC_w(v)$  to be the simple perverse sheaf generated by the rank-1 trivial local system on  $\mathcal{M}_0^{\bullet\text{reg}}(v, w)$ .

By the celebrated decomposition theorem [BBD82], the sheaf  $\pi_w(v) = \pi_!(1_{\mathcal{M}^\bullet(v, w)})$  decomposes into a direct sum of shifts of simple perverse sheaves on  $\mathcal{M}_0^\bullet(w)$ . The results in [Nak01a, Theorem 14.3.2] can be translated as follows.

**Theorem 5.2.1.** *We have a decomposition*

$$(44) \quad \pi_w(v) = \oplus_{v': (v', w) \text{ is } l\text{-dominant}} \oplus_{d \in \mathbb{Z}} a_{v, v'; w}^d IC_w(v')[d],$$

where the coefficients  $a_{v, v'; w}^d$  satisfy  $a_{v, v'; w}^d \in \mathbb{N}$ ,  $a_{v, v'; w}^d = a_{v, v'; w}^{-d}$ ,  $a_{v, v'; w}^d = \delta_{d0}$  if  $(v, w)$  is  $l$ -dominant, and  $a_{v, v'; w}^d$  vanishes unless  $(v', w) \geq (v, w)$ .

*Proof.* By Proposition 4.3.5, Proposition 4.3.12, Corollary 4.3.14, the summands appearing in the decomposition of  $\pi_!(1_{\mathcal{M}^\bullet(v, w)})$  are shifts of simple perverse sheaves generated by local systems over  $\mathcal{M}_0^{\bullet\text{reg}}(v', w)$  for  $l$ -dominant pairs  $(v', w)$ .

The argument in the proof of [Nak01a, Theorem 14.3.2] implies the following results:

- (1)  $a_{v, v'; w}^d = \delta_{d0}$  if  $(v, w)$  is  $l$ -dominant, since  $\pi$  is an isomorphism between  $\pi^{-1}\mathcal{M}_0^{\bullet\text{reg}}(v, w) \subset \mathcal{M}^\bullet(v, w)$  and  $\mathcal{M}_0^{\bullet\text{reg}}(v, w)$ ;
- (2) Because we have Theorem 4.3.4, the transversal slice technique in the proof of [Nak01a, Theorem 14.3.2] is effective. It follows that the decomposition summands are shifts of simple perverse sheaves  $IC_w(v')$  generated by trivial local systems.

Finally, since  $1_{\mathcal{M}^\bullet(v, w)}$  is invariant under the Verdier duality  $D$ , the sheaf  $\pi_w(v)$  is invariant as well, and therefore we have  $a_{v, v'; w}^d = a_{v, v'; w}^{-d}$ .  $\square$

Let  $a_{v, v'; w}(t)$  denote the Laurent polynomial in the variable  $t$  given by

$$a_{v, v'; w}(t) = \sum_{d \in \mathbb{Z}} a_{v, v'; w}^d t^d.$$

Theorem 4.3.4 implies that we have  $a_{v, v'; w}(t) = a_{v-v', 0; w-C_q v}(t)$ .

For each  $l$ -dominant pair  $(v, w)$ , we define the set

$$\mathcal{P}_w = \{IC_w(v) | (v, w) \text{ is } l\text{-dominant}, \}.$$

It is a finite set by Remark 4.3.13.

Let  $\mathcal{D}_c(\mathcal{M}_0^\bullet(w))$  be the bounded derived category of constructible sheaves on  $\mathcal{M}_0^\bullet(w)$ , and  $\mathcal{Q}_w$  its full subcategory whose objects are isomorphic to the direct sums of the shifts of the objects in  $\mathcal{P}_w$ . Then  $\mathcal{Q}_w$  and  $\mathcal{P}_w$  are stable under the Verdier duality  $D$ . Let  $\mathcal{K}_w$  be the quotient of the free abelian group generated by the isomorphism classes  $(L)$  in  $\mathcal{Q}_w$  modulo the relation

$(L) = (L') + (L'')$  whenever  $L$  is isomorphic to  $L' \oplus L''$ . The group  $\mathcal{K}_w$  has a natural  $\mathbb{Z}[t^\pm]$ -structure such that  $t(L) = (L[1])$ . The duality  $D$  induces an involution  $(\overline{\phantom{x}})$  on  $\mathcal{K}_w$  which satisfies  $t(\overline{L}) = t^{-1}(\overline{L})$  and  $(\overline{IC_w(v)}) = (IC_w(v))$ .

The decomposition (44) implies that, by abuse of notation, we obtain that  $\mathcal{K}_w$  has a  $\mathbb{Z}[t^\pm]$ -basis  $\{IC_w(v) | (v, w) \text{ is } l\text{-dominant}\}$  and a  $\mathbb{Z}[t^\pm]$ -basis  $\{\pi_w(v) | (v, w) \text{ is } l\text{-dominant}\}$ .

As in [Nak11], we define the abelian group  $\mathcal{K}_w^* = \text{Hom}_{\mathbb{Z}[t^\pm]}(\mathcal{K}_w, \mathbb{Z}[t^\pm])$ . Let  $\{L_w(v)\}$ ,  $\{\chi_w(v)\}$  be the bases of  $\mathcal{K}_w^*$  dual to  $\{IC_w(v)\}$ ,  $\{\pi_w(v)\}$  respectively. Define another basis  $\{M_w(v) | (v, w) \text{ is } l\text{-dominant}\}$  of  $\mathcal{K}_w^*$  by

$$\langle M_w(v), L \rangle = \sum_k t^{\dim \mathcal{M}_0^{\bullet \text{reg}}(v, w) - k} \dim H^k(i_{x_{v, w}}^! L),$$

where  $x_{v, w}$  is any point in  $\mathcal{M}_0^{\bullet \text{reg}}(v, w)$ ,  $i_{x_{v, w}}$  is the inclusion, and  $\langle \phantom{x}, \phantom{x} \rangle$  is the canonical pairing. Notice that the definition of  $M_w(v)$  is independent of the choice of  $x_{v, w}$ . Indeed, it suffices to check this for the elements  $L$  in the basis  $\{\pi_w(v)\}$  and here it follows from Theorem 4.3.4.

In the situation of Theorem 4.3.4, we have

$$\langle M_w(v'), IC_w(v) \rangle = \langle M_{w^\perp}(v'^\perp), IC_{w^\perp}(v^\perp) \rangle.$$

By the properties of perverse sheaves, we have

$$(45) \quad L_w(v) \in M_w(v) + \sum_{(v', w) < (v, w)} t^{-1} \mathbb{Z}[t^{-1}] M_w(v').$$

Therefore,  $\{M_w(v) | (v, w) \text{ is } l\text{-dominant}\}$  is a basis.

Following [Nak04, Section 8], let us denote the inverse transform by

$$(46) \quad M_w(v) = L_w(v) + \sum_{(v', w) < (v, w)} Z_{v, v'; w}(t) L_w(v'),$$

where  $Z_{v, v'; w}(t) \in t^{-1} \mathbb{Z}[t^{-1}]$ .

**Definition 5.2.2** ([Nak11, 3.3]). Define  $\mathcal{R}_t$  to be the infinite rank free  $\mathbb{Z}[t^\pm]$ -module consisting of the functionals  $(f_w) \in \prod_w \mathcal{K}_w^*$  such that  $\langle f_w, IC_w(v) \rangle$  equals  $\langle f_{w^\perp}, IC_{w^\perp}(v^\perp) \rangle$  for any  $l$ -dominant pairs  $(v, w)$ ,  $(w^\perp, v^\perp)$  appearing in Theorem 4.3.4.

Let  $M(w) = (f_{w'})_{w'}$  denote the functional determined by  $f_w = M_w(0)$  and  $L(w) = (f_w)_{w'}$  the functional determined by  $f_w = L_w(0)$ . Then  $\{M(w)\}$  and  $\{L(w)\}$  are two bases of  $\mathcal{R}_t$ .

By [VV03], resp. [Nak11, Section 3.5], for any  $w, w^1, w^2$ , such that  $w^1 + w^2 = w$ , we have a restriction functor

$$\widetilde{\text{Res}}_{w^1, w^2}^w : \mathcal{D}_c(\mathcal{M}_0^{\bullet}(w)) \rightarrow \mathcal{D}_c(\mathcal{M}_0^{\bullet}(w^1)) \times \mathcal{D}_c(\mathcal{M}_0^{\bullet}(w^2)).$$

Furthermore,  $\widetilde{\text{Res}}_{w^1, w^2}^w$  sends  $\pi_w(v)$  to

$$\oplus_{v^1 + v^2 = v} \pi_{w^1}(v^1) \boxtimes \pi_{w^2}(v^2) [d((v^2, w^2), (v^1, w^1)) - d((v^1, w^1), (v^2, w^2))],$$

where  $(v^1, w^1), (v^2, w^2)$  are not necessarily  $l$ -dominant.

We define  $\text{Res}^w = \sum_{w^1 + w^2 = w} \widetilde{\text{Res}}_{w^1, w^2}^w [-\mathcal{E}'(w^1, w^2)]$  for each  $w$ . Because these functors are compatible with Theorem 4.3.4, they induce a multiplication of  $\mathcal{R}_t$ , which we denote by  $\otimes$ .

The arguments of [VV03] imply the following result.

**Theorem 5.2.3.** *The structure constants of the multiplication  $\otimes$  with respect to the basis  $\{L(w)\}$  of  $\mathcal{R}_t$  are positive:*

$$(47) \quad L(w^1) \otimes L(w^2) = \sum_{w^3} b_{w^1, w^2}^{w^3}(t) L(w^3)$$

with  $b_{w^1, w^2}^{w^3}(t) \in \mathbb{N}[t^\pm]$ .

*Proof.* The structure constants are determined by those of the coalgebra structure on perverse sheaves, cf. [VV03, Section 5.1 Lemma 1(b)].  $\square$

**5.3. Rings of characters.** Define the ring of formal power series

$$(48) \quad \widehat{\mathcal{Y}} = \mathbb{Z}[t^\pm][[W_i(a), V_i(b)]_{i \in I, a \in \mathbb{Z}, b \in (\mathbb{Z} + \frac{1}{2})}],$$

where  $t, W_i(a), V_i(a)$  are indeterminates. We denote its product by  $\cdot$ , and often omit this notation.

Given vectors  $w, v$  as before, we let  $m(v, w)$  denote the monomial  $m = \prod_{i,a} W_i(a)^{w_i(a)} \prod_{i,b} V_i(b)^{v_i(b)}$ . Endow  $\widehat{\mathcal{Y}}$  with the twisted product  $*$  and the bar involution  $\overline{(\ )}$  such that for any two monomials  $m^1 = m(v^1, w^1), m^2 = m(v^2, w^2)$ , we have

$$(49) \quad \bar{t} = t^{-1}, \quad \overline{m^1} = m^1,$$

$$(50) \quad m^1 * m^2 = t^{-\tilde{d}'(m^1, m^2) + \tilde{d}(m^2, m^1)} m^1 m^2.$$

We follow the notation of Section 5.2. Moreover, as in Theorem 4.3.4, we write  $\mathbf{m}^\bullet_x(v, w)$  for the fibre  $\pi^{-1}(x)$  of a point  $x$  under  $\pi : \mathcal{M}^\bullet(v, w) \rightarrow \mathcal{M}_0^\bullet(v, w)$ . Notice that the homology of the fibre  $H_*(\mathbf{m}^\bullet_{x_{v,w}}(v', w), \mathbb{C})$  is isomorphic to  $H^{\dim \mathcal{M}^\bullet(v', w) - *} (i_{x_{v,w}}^! \pi_w(v')),$  cf. [CG97, 8.5.4]. Therefore, Theorem 4.3.4 implies that

$$\langle M(v, w), \pi_w(v') \rangle = \langle M(v^\perp, w^\perp), \pi_{w^\perp}(v' - v^0) \rangle.$$

**Definition 5.3.1** (*t-analogue of q-characters*). *For any given  $w$ , define  $\widehat{\chi}_{q,t}(\ )$  to be the  $\mathbb{Z}[t^\pm]$ -linear map from  $\mathcal{K}_w^*$  to  $\widehat{\mathcal{Y}}$  such that*

$$(51) \quad \widehat{\chi}_{q,t}(\ ) = \sum_v \langle \ , \pi_w(v) \rangle W^w V^v.$$

We can compute

$$\begin{aligned} \widehat{\chi}_{q,t}(L_w(v)) &= \sum_{v'} a_{v', v; w}(t) W^w V^{v'} \\ &= \sum_{v' - v} a_{v' - v, 0; w}(t) W^w V^{v'}. \end{aligned}$$

Since  $W^w V^{v'}$  and  $a_{v',v;w}(t)$  are bar-invariant,  $\widehat{\chi_{q,t}}(\ )$  is bar-invariant as well. Similarly, we have

$$\begin{aligned}
M_w^{\widehat{\mathcal{Y}}}(v) &= \widehat{\chi_{q,t}}(M_w(v)) \\
&= \sum_{v',k} t^{\dim \mathcal{M}^\bullet(v,w)-k} H^k(i_{x_{v,w}}^! \pi_w(v'), \mathbb{C}) W^w V^{v'} \\
&= \sum_{v',k} t^{\dim \mathcal{M}^\bullet(v,w)-\dim \mathcal{M}^\bullet(v',w)} t^{\dim \mathcal{M}^\bullet(v',w)-k} H^k(i_{x_{v,w}}^! \pi_w(v'), \mathbb{C}) W^w V^{v'} \\
&= \sum_{v',l} t^{\dim \mathcal{M}^\bullet(v,w)-\dim \mathcal{M}^\bullet(v',w)} t^l H_l(\mathfrak{m}_{x_{v,w}}^\bullet(v', w), \mathbb{C}) W^w V^{v'} \\
&\stackrel{4.3.19}{=} \sum_{v'} t^{\dim \mathcal{M}^\bullet(v,w)-\dim \mathcal{M}^\bullet(v',w)} P_t(\mathfrak{m}_{x_{v,w}}^\bullet(v', w)) W^w V^{v'} \\
&= \sum_{v'-v} t^{-\dim \mathcal{M}^\bullet(v'-v, w-C_q v)} P_t(\mathcal{L}^\bullet(v'-v, w-C_q v)) W^w V^{v'}.
\end{aligned}$$

Here  $P_t(\ )$  is the Poincaré polynomial  $\sum_k (-t)^k \dim H_k(\ )$ .

**Definition 5.3.2** (dual PBW elements). *For any given  $w$ , define  $M_w^{\widehat{\mathcal{Y}}}(w)$  to be the generating series*

$$(52) \quad M_w^{\widehat{\mathcal{Y}}}(w) = \widehat{\chi_{q,t}}(M_w(0)) = \sum_v P_t(\mathcal{L}^\bullet(v, w)) t^{-d((v,w), (v,w))} W^w V^v.$$

Notice that the summation runs over all  $v$  and  $(v, w)$  is not necessarily  $l$ -dominant.

**Remark 5.3.3.** *By Theorem 4.3.19, the parameter  $-t$  in  $P_t(\ )$  in Definition 5.3.2 can be replaced by  $t$ .*

Let  $E(\ ; x, y) \in \mathbb{Z}[x, y]$  be the virtual Hodge Polynomial, cf. e.g. [HV08]. Define  $p_t(\ )$  to be the virtual Poincaré polynomial (called the  $E$ -polynomial in [Qin10]), i.e.  $p_t(\mathcal{L}^\bullet(v, w)) = E(\mathcal{L}^\bullet(v, w); t, t)$ . It is additive with respect to  $\alpha$ -partitions in the sense of [Nak01a, Section 7.1], and multiplicative with respect to fibre products.

**Proposition 5.3.4** ([Nak04, Lemma 5.2]). *We have  $p_t(\mathcal{L}^\bullet(v, w)) = P_t(\mathcal{L}^\bullet(v, w))$  and  $p_t(\mathcal{M}^\bullet(v, w)) = P_t(\mathcal{M}^\bullet(v, w))$ .*

Recall that, in  $\mathcal{K}_w^*$ , we have  $M_w(v) = \sum_{v'} Z_{(v,w)(v',w)}(t) L_w(v')$ , where the matrix of coefficients  $(Z_{(v,w)(v',w)})$  is an upper uni-triangular matrix with respect to the dominance order. Therefore, we can solve the following equation recursively

$$\overline{M_w(v)} = \sum_{v': (v',w) \leq (v,w)} u_{(v,w),(v',w)} M_w(v'),$$

where the coefficients  $u_{(v,w),(v',w)} \in \mathbb{Z}[t^\pm]$ .

Finally, we apply  $\widehat{\chi_{q,t}}$  to the above equation and obtain

$$(53) \quad \overline{M_w^{\widehat{\mathcal{Y}}}(v)} = \sum_{v': (v',w) \leq (v,w)} u_{(v,w),(v',w)} M_w^{\widehat{\mathcal{Y}}}(v'),$$

**Remark 5.3.5.** *In fact, we can compute the coefficient matrix  $(u_{(v,w)(v',w)})$  and the basis transition matrix  $(Z_{(v,w)(v',w)})$  from  $P_t(\mathcal{L}^\bullet(v,w))$ , cf. [Nak04, Section 8].*

**Proposition 5.3.6** ([Nak04, Proposition 6.2]). *If for all  $i, j \in I$ ,  $a, b \in \mathbb{Z}$ ,  $a > b$ , either  $(w^1)_i(a)$  or  $(w^2)_j(b)$  vanishes, then we have*

$$(54) \quad M^{\hat{\mathcal{Y}}}(w^2) * M^{\hat{\mathcal{Y}}}(w^1) = t^{\mathcal{E}'(w^1, w^2)} M^{\hat{\mathcal{Y}}}(w^1 + w^2).$$

*Proof.* As in the proof of [Nak04, Proposition 6.2], for any monomials  $m^1 = W^{w^1} V^{v^1}$ ,  $m^2 = W^{w^2} V^{v^2}$ , we have a complex vector bundle  $\widetilde{\mathcal{Z}}^\bullet(v^1, w^1; v^2, w^2)$  of rank  $\text{Ker } \tau^{21} / \text{Im } \sigma^{21} = d((v^1, w^1), (v^2, w^2))$  over each  $\mathcal{L}^\bullet(v^1, w^1) \times \mathcal{L}^\bullet(v^2, w^2)$ . The set of  $\widetilde{\mathcal{Z}}^\bullet(v^1, w^1; v^2, w^2)$  for various  $v^1, v^2$ , gives a stratification of  $\mathcal{L}^\bullet(v^1 + v^2, w^1 + w^2)$ , cf. [Nak01b, 6.12]. Using the additivity and multiplicativity of  $p_t(\cdot)$ , we deduce

$$\begin{aligned} & M^{\hat{\mathcal{Y}}}(w^2) * M^{\hat{\mathcal{Y}}}(w^1) \\ &= \sum_{v^1, v^2} p_t(\mathcal{L}^\bullet(v^1, w^1)) \times p_t(\mathcal{L}^\bullet(v^2, w^2)) t^{-d(m^1, m^1) - d(m^2, m^2)} m^2 * m^1 \\ &= \sum_{v^1, v^2} p_t(\mathcal{L}^\bullet(v^1, w^1) \times \mathcal{L}^\bullet(v^2, w^2)) t^{-d(m^1, m^1) - d(m^2, m^2)} \\ & \quad t^{\tilde{d}'(m^1, m^2) - \tilde{d}'(m^2, m^1)} m^1 m^2 \\ &= \sum_{v^1, v^2} p_t(\widetilde{\mathcal{Z}}^\bullet(v^1, w^1; v^2, w^2)) t^{-2d(m^1, m^2)} t^{-d(m^1, m^1) - d(m^2, m^2)} \\ & \quad t^{\mathcal{E}'(w^1, w^2) + d(m^1, m^2) - d(m^2, m^1)} m^1 m^2 \\ &= \sum_{v^1, v^2} p_t(\widetilde{\mathcal{Z}}^\bullet(v^1, w^1; v^2, w^2)) t^{\mathcal{E}'(w^1, w^2)} t^{-d(m^1 m^2, m^1 m^2)} m^1 m^2 \\ &= \text{RHS}. \end{aligned}$$

□

Define the ring of formal power series

$$(55) \quad \mathcal{Y} = \mathbb{Z}[t^\pm][[Y_i(a)^\pm]]_{i \in I, a \in \mathbb{Z}},$$

where  $t, Y_i(a)$  are indeterminates. We denote its product by  $\cdot$ , and often omit this notation.

Given vectors  $w, v$  as before, we denote the monomial  $m_Y = Y^{w - C_q v}$  by  $m_Y(v, w)$ . By (36), we have a naturally defined bilinear form  $\tilde{d}'$  for such monomials. For  $i \in I$ ,  $b \in (\frac{1}{2} + \mathbb{Z})$ , we sometimes denote  $m_Y(e_{i,b}, 0)^{-1}$  by  $A_{i,b}$ .

Endow  $\mathcal{Y}$  with the twisted product  $*$  and the bar involution  $\overline{(\cdot)}$  such that for any two monomials  $m_Y^1 = m_Y(v^1, w^1)$ ,  $m_Y^2 = m_Y(v^2, w^2)$ , we have

$$(56) \quad \bar{t} = t^{-1}, \quad \overline{m_Y^1} = m_Y^1,$$

$$(57) \quad m_Y^1 * m_Y^2 = t^{-\tilde{d}'(m_Y^1, m_Y^2) + \tilde{d}'(m_Y^2, m_Y^1)} m_Y^1 m_Y^2.$$

Define the  $\mathbb{Z}[t^\pm]$ -linear map  $\hat{\Pi}$  from  $\hat{\mathcal{Y}}$  to  $\mathcal{Y}$  such that for any  $(v, w)$ , it sends  $m(v, w)$  to  $m_Y(v, w)$ . Then it is a ring homomorphism. The  $t$ -analogue

of the  $q$ -character map is defined to be the  $\mathbb{Z}[t^\pm]$ -linear map  $\chi_{q,t}(\ )$  from  $\mathcal{R}_t$  to  $\mathcal{Y}$  such that we have

$$(58) \quad \chi_{q,t}(\ ) = \sum_v \langle \ , \pi(v, w) \rangle Y^{w - C_q v}.$$

The following results follow from [VV03] and [Nak04, Theorem 3.5].

**Theorem 5.3.7.** 1)  $\widehat{\chi_{q,t}}(\ )$  is injective.

2)  $\chi_{q,t}(\ )$  is an injective algebra homomorphism from  $\mathcal{R}_t$  to  $\mathcal{Y}$ .

## 6. RESULTS ON GENERIC CHARACTERS

Following Nakajima's work [Nak11], we could find a link between truncated  $qt$ -characters and cluster characters by studying certain Fourier-Deligne-Sato transform. This approach is the key ingredient of the first part of the paper [KQ12]. In this section, we recall useful results from [KQ12].

Let us fix the following conventions.

- We always assume  $w \in \mathbb{N}^{I \times \{-1, 0\}}$  (level 1 case). Furthermore, we assume  $v \in \mathbb{N}^{I \times \{-\frac{1}{2}\}}$ , which is naturally identified with a dimension vector  $v \in \mathbb{N}^I$ .
- All the representations are those of the opposite quiver  $Q^{op}$ .
- We assume the ice quiver  $\tilde{Q}$  is of level 1 with  $z$ -pattern.

Since our main interest lies in quantum cluster algebras, we only study the *truncated  $qt$ -characters* in the sense of [HL10]. By putting the above restrictions on the definitions of  $\widehat{\chi_{q,t}}$  and  $\chi_{q,t}$ , we obtain the truncated character maps  $\widehat{\chi_{q,t}}^{\leq 0}$  and  $\chi_{q,t}^{\leq 0}$ . Notice that the truncation preserves all the  $l$ -dominant pairs. Proposition 5.3.6 and Theorem 5.3.7 still hold for the truncated characters, cf. [HL10, Proposition 6.1].

**6.1. Fourier-Deligne-Sato transform.** For each multiplicity parameter  $m = (m_i)_{i \in I} \in \mathbb{N}^I$ , define the injective  $Q^{op}$ -representation  $I^m = \oplus_{i \in I} I_i^{\oplus m_i}$  and similarly the projective representation  $P^m = \oplus_{i \in I} P_i^{\oplus m_i}$ .

Let  $(B, \alpha, \beta)$  be any point of  $\text{Rep}^\bullet(v, w)$  and  $r$  any path of  $Q^{op}$ . Define<sup>5</sup> the homomorphism  $z_r$  to be the composition  $\beta_{t(r)} B_r \alpha_{s(r)}$  along the path  $\beta_{t(r)} r \alpha_{s(r)}$  if  $r$  is nontrivial and  $\beta_{t(r)} \alpha_{s(r)}$  if  $r$  is  $e_i$  for some vertex  $i \in I$ . Notice that each  $z_r$  determines a morphism from  $P_{s(r)}^{w_{s(r)}(0)}$  to  $I_{t(r)}^{w_{t(r)}(-1)}$ . Furthermore, we have the following result.

**Proposition 6.1.1** ([KQ12]). *When  $v$  is big enough, the affine quiver variety  $\mathcal{M}_0^\bullet(v, w)$  stabilizes to the affine space  $E_w = \text{Hom}(P^{w(0)}, I^{w(-1)})$ .*

**Definition 6.1.2.** *For each dimension vector  $v \in \mathbb{N}^I$ , let  $\mathcal{F}(v, w)$  denote the variety parameterizing all the  $I$ -graded submodules  $X = (X_i)_{i \in I}$  of  $I^{w(-1)}$ , such that  $\dim X = v$ .*

Since  $Q^{op}$  is acyclic, the variety  $\mathcal{F}(v, w)$  is smooth projective and irreducible [Rei08, Theorem 4.10].

<sup>5</sup> In [Nak11], our  $z_r$  is denoted by  $y_r$  if  $r$  is nontrivial and  $x_i$  if  $r$  is  $e_i$  for some vertex  $i \in I$ .

We define the variety  $\tilde{\mathcal{F}}(v, w)$  by

$$(59) \quad \tilde{\mathcal{F}}(v, w) = \{((X, z) \in \mathcal{F}(v, w) \times E_w \mid \text{Im } z \subset X)\}.$$

**Proposition 6.1.3** ([KQ12]). *The quasi-projective quiver variety  $\mathcal{M}^\bullet(v, w)$  is isomorphic to the variety  $\tilde{\mathcal{F}}(v, w)$ .*

Let  $E_w^*$  denote the natural dual space of the complex vector space  $E_w$ . Let us write  $\tilde{\mathcal{F}}(v, w)$  for the annihilator sub-bundle in  $\mathcal{F}(v, w) \times E_w^*$  of the sub-bundle  $\tilde{\mathcal{F}}(v, w) \subset \mathcal{F}(v, w) \times E_w$ . Its fibre over any given point  $X = (X_i) \in \mathcal{F}(v, w)$  consists of the linear maps  $z^* = (z_r^*) \in E_w^*$  such that, for any point  $(X, z) \in \tilde{\mathcal{F}}(v, w)$ , the natural pairing  $\langle z, z^* \rangle = \sum_r \text{Tr}(z_r^* z_r)$  vanishes.

Using the Nakayama functor  $\nu(\cdot)$ , we obtain

$$E_w^* = (\text{Hom}(P^{w(0)}, I^{w(-1)}))^* = \text{Hom}(I^{w(-1)}, \nu(P^{w(0)})) = \text{Hom}(I^{w(-1)}, I^{w(0)}).$$

**Lemma 6.1.4** ([KQ12]). *The fibre of  $\tilde{\mathcal{F}}(v, w)^\perp$  over any given point  $X \in \mathcal{F}(v, w)$  consists of the maps  $z^* = (z_r^*) \in \text{Hom}(I^{w(-1)}, I^{w(0)})$  such that*

$$(60) \quad z^* X = 0$$

Fix any element  $z^* \in \text{Hom}(I^{w(-1)}, I^{w(0)})$ . We define  ${}^\sigma W = ({}^\sigma W_i)_{i \in I}$  to be the kernel of  $z^*$  and denote its dimension vector by  ${}^\sigma w$ . Then  $({}^\sigma W, z^*)$  is contained in  $\tilde{\mathcal{F}}^\perp({}^\sigma w, w)$ . For any  $1 \leq i \leq n$ ,  $X_i$  is a subspace of  ${}^\sigma W_i$ .

**Definition 6.1.5.** *For any vector  $w \in \mathbb{N}^{I \times \{-1, 0\}}$ , its coefficient-free part  $\phi_w \in \mathbb{N}^{I \times \{-1, 0\}}$  and its pure coefficient part  ${}^f w \in \text{span}_{\mathbb{N}}\{e_{i, -1} + e_{i, 0} \mid i \in I\}$  are defined such that  $w = \phi_w + {}^f w$  and, for any  $i \in I$ , either  $\phi_{w_i}(-1)$  or  $\phi_{w_i}(0)$  vanishes. Let  $\mathcal{J}$  denote the set of  $w$  such that  $w = \phi_w$ .*

**Proposition 6.1.6** ([KQ12]). *1) The fibre of  $\tilde{\mathcal{F}}^\perp(v, w)$  over  $z^*$  is isomorphic to the submodule Grassmannian consisting of the  $v$ -dimensional submodules of  ${}^\sigma W$ .*

*2) When  $z^*$  is generic, the module  ${}^\sigma W$  is a generic representation of  $Q^{op}$  with dimension  ${}^\sigma w$ .*

**6.2. Generic characters.** Let the Fourier-Deligne-Sato transform from  $\pi : \tilde{\mathcal{F}}(v, w) \rightarrow E_w$  to  $\pi^\perp : \tilde{\mathcal{F}}^\perp(v, w) \rightarrow E_w^*$  be denoted by  $\Psi$ . Define the set

$$\mathcal{L}_w = \{IC_w(v) \in \mathcal{P}_w \mid \text{supp } \Psi IC_w(v) = E_w^*\}.$$

Recall that  $\Psi(IC_w(v))$  is a perverse sheaf over  $E_w^*$ .

**Definition 6.2.1** (Twisted rank). *For any  $w, w' \in \mathbb{N}^{I \times \{-1, 0\}}$ , we define the integer  $r_{w, w'}$  to be*

$$(61) \quad r_{w, w'} = r(v, w) = (-1)^{\dim_{\mathbb{C}} \mathcal{M}_0^\bullet(v, w)} \text{rank } \Psi(IC_w(v))$$

*if we have  $w' = w - C_q v$  for some  $v \in \mathbb{N}^{I \times \{\frac{1}{2}\}}$  such that  $\text{supp } \Psi(IC_w(v)) = E_w^*$  and zero otherwise.*

Notice that  $IC(0, w)$  is always contained in  $\mathcal{L}_w$ . It follows that  $r_{w, w} = 1$ . Furthermore, for each fixed  $w$ , only finitely many of the  $r_{w, w'}$  are nonzero. The matrix  $(r_{w, w'})$  is upper unitriangular with respect to the dominance order.

**Remark 6.2.2.** Our definition of  $r_{w,w'}$  is the twisted version of that of [Nak11, 6.1].

We define the *almost simple pseudo-module*  $\mathbb{L}(w)$  to be the element in the Grothendieck group  $\mathcal{R}_t$  such that we have

$$(62) \quad \mathbb{L}(w) = \sum_{w'} r_{w,w'} L(w').$$

Denote the truncated  $qt$ -characters  $\chi_{q,t}^{\leq 0}(\mathbb{L}(w))$  by  $\mathbb{L}^{\mathcal{Y}}(w)$ .

The following is the main theorem of the first part of [KQ12].

**Theorem 6.2.3** ([KQ12]). *The truncated  $qt$ -characters of the almost simple pseudo-modules in  $\mathcal{Y}$  are given by*

$$(63) \quad \mathbb{L}^{\mathcal{Y}}(w) = \sum_v t^{-\dim(\mathrm{Gr}_v^\sigma W)} P_t(\mathrm{Gr}_v^\sigma W) Y^{w-C_q v}.$$

Furthermore, the truncated characters of almost simple pseudo-modules have the following properties.

**Proposition 6.2.4** ([KQ12]). *1) If  $M$  is a rigid module with a minimal injective resolution  $z^* \in E_w$ , then  $\mathbb{L}(w) = L(w)$ .*

*2) If a generic map  $z^*$  in  $E_w^*$  has a rigid kernel, then  $\mathbb{L}(w) = L(w)$ .*

*3) In the notation of Definition 6.1.5, we have a factorization*

$$(64) \quad \mathbb{L}^{\mathcal{Y}}(w) = \mathbb{L}^{\mathcal{Y}}(\phi w) \cdot \mathbb{L}^{\mathcal{Y}}(f w).$$

Therefore, we also call them the *generic characters*.

## 7. FROM CHARACTERS TO THE QUANTUM TORUS

In this section, we construct  $\mathbb{Z}$ -linear maps from the ring of formal power series  $\widehat{\mathcal{Y}}$  in which  $qt$ -characters live to the quantum tori  $D(\mathcal{T})$  and  $\mathcal{T}$  in which quantum cluster algebras live. Although the maps fail to preserve some properties of the products of  $qt$ -characters, we find that the failures are bearable and controllable. This observation will play a crucial role in later sections.

**7.1. Dual PBW basis elements.** As in Section 2.2, let  $\widetilde{Q}$  be an ice quiver whose principal part  $Q$  is acyclic. Using the notation of Section 2.3, we define the linear map  $\mathrm{ind}(\cdot)$  from the set of finitely supported vectors in  $\mathbb{N}^{I \times \mathbb{Z}}$  to  $\mathbb{Z}^m$  such that  $\mathrm{ind}(e_{i,a})$ ,  $i \in I$ ,  $a \in \mathbb{Z}$ , is the vector of coordinates in the basis  $[e_i \Gamma]$ ,  $1 \leq i \leq m$ , of the index of the coefficient-free object whose image in the cluster category  $\mathcal{C}_Q$  is  $T_i[-a]$ .

**Lemma 7.1.1.** *We have, for  $1 \leq k \leq n$ ,*

$$(65) \quad \mathrm{ind}^z(e_{k,0}) = e_k,$$

$$(66) \quad \mathrm{ind}^z(e_{k,-1}) = e_{k+n} - e_k.$$

*Proof.* The first equation is clear.

Consider the triangle

$$T_k \xrightarrow{f} T_{k+n} \rightarrow M \rightarrow \Sigma T_k$$

in  $\mathcal{C}_{(\tilde{Q}, \tilde{W})}$ , where  $f$  is non-zero. Then  $f$  is a left approximation of  $T_k$  in  $\text{add}(T_{n+i}, 1 \leq i \leq n)$  in the sense of [IY08]. It follows that  $M$  is the coefficient-free object lifting  $T_i[1]$  and so we have  $\text{ind}^z(e_{k,-1}) = \text{ind } M = e_{k+n} - e_k$ .  $\square$

Let  $\text{pr}_n$  denote the projection of  $\mathbb{Z}^m$  (respectively  $\mathbb{Z}^{2n}$ ) to the first  $n$  coordinates. Then  $\text{pr}_n \text{ind}$  is independent of the choice of the coefficient  $\tilde{Q} - Q$ .

**Lemma 7.1.2.** *We have*

$$(67) \quad \text{ind}^z(w - C_q v) = \text{ind}^z(w) + \tilde{B}^z v.$$

*Proof.* It suffices to check the case of  $w = 0$  and  $v = e_{k,-\frac{1}{2}}$ ,  $1 \leq k \leq n$ . We have

$$\begin{aligned} \text{ind}^z(-C_q v) &= -\text{ind}^z(e_{k,-1} + e_{k,0}) + \sum_{i:1 \leq i < k} b_{ik} \text{ind}^z(e_{i,0}) \\ &\quad + \sum_{j:k < j \leq n} b_{kj} \text{ind}^z(e_{j,-1}) \\ &= -e_{k+n} + \sum_{1 \leq i < k} b_{ik} e_i + \sum_{k < j \leq n} b_{kj} (e_{j+n} - e_j) \\ &= b_{k+n,k} e_{k+n} + \sum_{1 \leq i < k} b_{ik} e_i + \sum_{k < j \leq n} b_{j+n,k} e_{j+n} + \sum_{k < j \leq n} b_{jk} e_j \\ &= \tilde{B}^z e_k. \end{aligned}$$

$\square$

Let  $\delta$  be a strictly positive integer. Denote the product of  $\delta$  with the rank  $m$  identity matrix  $I_m$  by  $D$ . Let  $(\tilde{B}, \Lambda)$  be a compatible pair such that  $\Lambda(-\tilde{B}) = \begin{bmatrix} D \\ 0 \end{bmatrix}$ .

Define  $\tilde{N} = \max\{2n, m\}$ . Extend  $\Lambda$  into an  $N \times N$  matrix by putting zero entries in the extra rows and columns indexed by either  $\{m+1, \dots, N\}$  or  $\{n+1, \dots, N\}$ . We extend the  $m \times n$  matrix  $\tilde{B}$  and the  $2n \times n$  matrix  $\tilde{B}^z$  to  $N \times N$  matrices similarly. Notice that  $\Lambda(e_i, \tilde{B}v) = 0$ , for any  $n+1 \leq i \leq N$  and  $v \in \mathbb{N}^n$ .

**Definition 7.1.3.** *The double quantum torus is the Laurent polynomial ring*

$$(68) \quad D(\mathcal{T}) = \mathbb{Z}[q^{\pm \frac{1}{2}}][x_1^{\pm}, \dots, x_N^{\pm}, y_1^{\pm}, \dots, y_n^{\pm}],$$

together with the twisted product  $*$  such that for any  $g^1, g^2 \in \mathbb{Z}^N$ ,  $v^1, v^2 \in \mathbb{Z}^n$ , we have

$$x^{g^1} y^{v^1} * x^{g^2} y^{v^2} = q^{\frac{1}{2} \Lambda(g^1 + \tilde{B}v^1, g^2 + \tilde{B}v^2)} x^{g^1 + g^2} y^{v^1 + v^2}.$$

It has the bar involution  $\overline{(\ )}$  given by  $\overline{(q^{\frac{1}{2}} x^g y^v)} = q^{-\frac{1}{2}} x^g y^v$ .

Define the quantum torus  $\mathcal{T}$  to be the  $\mathbb{Z}[q^{\pm\frac{1}{2}}]$ -subalgebra of  $D(\mathcal{T})(+, *)$  generated by  $x_1^\pm, \dots, x_N^\pm$ . The coefficient ring is  $\mathbb{Z}P[q^{\pm\frac{1}{2}}] = \mathbb{Z}[q^{\pm\frac{1}{2}}][x_{n+1}^\pm, \dots, x_m^\pm]$ . Let  $\mathbb{Z}P$  denote its semi-classical limit under the specialization  $q^{\frac{1}{2}} \mapsto 1$ .

Next, we construct a map from  $\widehat{\mathcal{Y}}$  to  $D(\mathcal{T})$ .

**Definition 7.1.4** (Correspondence map). *The  $\mathbb{Z}$ -linear map  $\text{cor}$  from  $\widehat{\mathcal{Y}}$  to  $D(\mathcal{T})$  is given by*

$$(69) \quad \text{cor}(t^\lambda W^w V^v) = q^{\frac{\delta}{2}\lambda} x^{\text{ind}(w)} y^v,$$

for any  $w, v$ , and integer  $\lambda$ .

Notice that  $\text{cor}$  commutes with the bar involutions. Although  $\text{cor}$  does not commute with twisted products  $*$ , we can measure the failure as below.

**Lemma 7.1.5** (Failure 1). *We have, for any monomials  $m^i = W^{w^i} V^{v^i}$ ,  $i = 1, 2$ ,*

$$(70) \quad \begin{aligned} \text{cor}(W^{w^1} V^{v^1} * W^{w^2} V^{v^2}) &= q^{\frac{\delta}{2}\Lambda^z(\text{ind}^z(w^1), \text{ind}^z(w^2)) - \frac{1}{2}\Lambda(\text{ind}(w^1), \text{ind}(w^2))} \\ &\quad \text{cor}(W^{w^1} V^{v^1}) * \text{cor}(W^{w^2} V^{v^2}). \end{aligned}$$

In particular, the  $q$ -power does not depend on  $v^1, v^2$ .

*Proof.* Notice that we always have  $\Lambda(\text{ind } w, \widetilde{B}v) = -\delta \text{pr}_n \text{ind } w \cdot v$ , and  $\Lambda(\widetilde{B}v^1, \widetilde{B}v^2) = \delta \langle v^1, v^2 \rangle - \delta \langle v^2, v^1 \rangle$ , cf. [Qin10, 5.2.1]. Using Lemma 5.1.3 and 7.1.2, we obtain

$$\begin{aligned} \text{LHS} &= \text{cor}(t^{-\widetilde{d}'(m^1, m^2) + \widetilde{d}'(m^2, m^1)} m^1 m^2) \\ &= \text{cor}(t^{-\mathcal{E}'(w^1 - C_q v^1, w^2 - C_q v^2)} m^1 m^2) \\ &= q^{\frac{\delta}{2}\Lambda^z(\text{ind}^z(w^1 - C_q v^1), \text{ind}^z(w^2 - C_q v^2))} X^{\text{ind } w^1} Y^{v^1} X^{\text{ind } w^2} Y^{v^2} \\ &= q^{\frac{\delta}{2}\Lambda^z(\text{ind}^z w^1 + \widetilde{B}^z v_1, \text{ind}^z w^2 + \widetilde{B}^z v_2)} q^{-\frac{1}{2}\Lambda(\text{ind } w^1 + \widetilde{B}v^1, \text{ind } w^2 + \widetilde{B}v^2)} \\ &\quad X^{\text{ind } w^1} Y^{v^1} * X^{\text{ind } w^2} Y^{v^2} \\ &= q^{\frac{\delta}{2}\Lambda^z(\text{ind}^z w^1, \text{ind}^z w^2)} q^{-\frac{1}{2}\Lambda(\text{ind } w^1, \text{ind } w^2)} X^{\text{ind } w^1} Y^{v^1} * X^{\text{ind } w^2} Y^{v^2} \\ &= \text{RHS}. \end{aligned}$$

□

**Definition 7.1.6.** *We define the  $\mathbb{Z}[q^{\pm\frac{1}{2}}]$ -linear map  $\widehat{\Pi}$  from  $D(\mathcal{T})$  to  $\mathcal{T}$  such that it sends  $x^g y^v$  to  $x^{g+\widetilde{B}v}$ .*

Notice that  $\widehat{\Pi}$  is an algebra homomorphism with respect to both the usual products and the twisted products.

Let us define

$$(71) \quad M^{D(\mathcal{T})}_w(v) = \text{cor} \widehat{\chi_{q,t}}^{\leq 0}(M_w(v)),$$

$$(72) \quad M^{\mathcal{T}}_w(v) = \widehat{\Pi} M^{D(\mathcal{T})}_w((v)).$$

When  $v = 0$ , we also denote them by  $M^{D(\mathcal{T})}(w)$  and  $M^{\mathcal{T}}(w)$ . Explicitly, we have

$$\begin{aligned} M^{D(\mathcal{T})}(w) &= \sum_v P_{q^{\frac{\delta}{2}}}(\mathcal{L}(v, w)) q^{-\frac{\delta}{2} \dim \mathcal{M}^\bullet(v, w)} x^{\text{ind}(w)} y^v \\ &= \sum_v \text{cor}(\langle M_w(0), \pi_w(v) \rangle) x^{\text{ind}(w)} y^v. \end{aligned}$$

Recall that we have a similar map from  $\widehat{\mathcal{Y}}$  to  $\mathcal{Y}$ , which is also denoted by  $\widehat{\Pi}$ . In general, we do *not* have  $\widehat{\Pi} \widehat{\chi_{q,t}} = \chi_{q,t} \widehat{\Pi}$ . This failure is measured by the following result.

**Lemma 7.1.7** (Failure 2). *We have*

$$(73) \quad M^{\mathcal{T}}_w(v) = M^{\mathcal{T}}(w - C_q v) x^{\widetilde{B}v + \text{ind } C_q v}.$$

*Proof.* Straightforward calculation shows

$$\begin{aligned} M^{\mathcal{T}}_w(v) &= \sum_{v': v' - v \in \mathbb{N}^I} \widehat{\Pi} \text{cor}(\langle M_w(v), \pi_w(v') \rangle) x^{\text{ind}(w)} x^{\widetilde{B}v'}. \\ &= \sum_{v' - v} \widehat{\Pi} \text{cor}(\langle M_{w - C_q v}(0), \pi_w(v' - v) \rangle) x^{\text{ind}(w - C_q v)} x^{\widetilde{B}(v' - v)} \\ &\quad x^{\widetilde{B}v + \text{ind}(w) - \text{ind}(w - C_q v)}. \\ &= M^{\mathcal{T}}(w - C_q v) x^{\widetilde{B}v + \text{ind } C_q v}. \end{aligned}$$

□

Notice that the *correction factor*  $x^{\widetilde{B}v + \text{ind } C_q v}$  is contained in  $\mathbb{Z}P[q^{\pm \frac{1}{2}}]$ .

**Proposition 7.1.8.** *We have*

$$(74) \quad \overline{M^{\mathcal{T}}(w)} = \sum_{(v, w) \leq (0, w)} u_{(0, w), (v, w)} x^{\widetilde{B}v + \text{ind } C_q v} M^{\mathcal{T}}(w - C_q v).$$

*Proof.* If we apply  $\widehat{\Pi} \text{cor}$  to (53), the statement follows from (73). □

**Proposition 7.1.9.** *Fix  $w^1$  and  $w^2$ . If for all  $i, j \in I$  and  $a > b \in \mathbb{Z}$ , either  $(w^1)_i(a)$  or  $(w^2)_j(b)$  vanishes, then the multiplicative property holds:*

$$\begin{aligned} M^{\mathcal{T}}(w^2) * M^{\mathcal{T}}(w^1) &= q^{\frac{1}{2} \Lambda(\text{ind}(w^2), \text{ind}(w^1)) - \frac{\delta}{2} \Lambda^z(\text{ind}^z(w^2), \text{ind}^z(w^1))} \\ &\quad q^{\frac{\delta}{2} \mathcal{E}'(w^1, w^2)} M^{\mathcal{T}}(w^1 + w^2). \end{aligned}$$

*Proof.* Using (70), (73), and Proposition 5.3.6, we obtain

$$\begin{aligned} M^{\mathcal{T}}(w^2) * M^{\mathcal{T}}(w^1) &= \widehat{\Pi}(\text{cor} \widehat{\chi_{q,t}}^{\leq 0}(M(w^2)) * \text{cor} \widehat{\chi_{q,t}}^{\leq 0}(M(w^1))) \\ &= q^{\frac{1}{2} \Lambda(\text{ind}(w^2), \text{ind}(w^1)) - \frac{\delta}{2} \Lambda^z(\text{ind}^z(w^2), \text{ind}^z(w^1))} \\ &\quad \widehat{\Pi} \text{cor}(\widehat{\chi_{q,t}}^{\leq 0}(M(w^2)) * \widehat{\chi_{q,t}}^{\leq 0}(M(w^1))) \\ &= \text{RHS}. \end{aligned}$$

□

Let  $\mathcal{A}_{\text{sub}}^q$  denote the vector space spanned by the standard basis elements  $M^{\mathcal{T}}(w)$  over  $\mathbb{Z}P[q^{\pm \frac{1}{2}}]$ . The following is the main result of this section.

**Proposition 7.1.10.** *The vector space  $\mathcal{A}_{sub}^q$  is closed under the involution  $\overline{(\ )}$  and the twisted multiplication  $*$ .*

*Proof.* The first assertion follows from Proposition 7.1.8. It remains to verify the second one.

Recall that  $w_i(a)$  is the  $(i, a)$ -th component of  $w$ , and  $w(a) = \oplus_i w_i(a)$ , where  $i \in I$ ,  $a \in \{-1, 0\}$ . We have  $M(w_i(a)) = L(w_i(a))$ . Consequently,  $M^\mathcal{T}(w_i(a)) = \widehat{\Pi\text{cor}}_{\chi_{q,t}}^{\leq 0}(L(w_i(a)))$  is bar-invariant. Applying Proposition 7.1.9, we obtain that  $M^\mathcal{T}(w(a))$  is bar-invariant.

Use Propositions 7.1.8 and 7.1.9. For any two elements  $M^\mathcal{T}(w^1)$ ,  $M^\mathcal{T}(w^2)$ , up to specified invertible elements in  $\mathbb{Z}[q^{\pm \frac{1}{2}}]$ ,  $M^\mathcal{T}(w^1) * M^\mathcal{T}(w^2)$  becomes

$$\begin{aligned} & M^\mathcal{T}(w^1(0)) * M^\mathcal{T}(w^1(-1)) * M^\mathcal{T}(w^2(0)) * M^\mathcal{T}(w^2(-1)) \\ &= M^\mathcal{T}(w^1(0)) * \overline{M^\mathcal{T}(w^2(0))} * \overline{M^\mathcal{T}(w^1(-1))} * M^\mathcal{T}(w^2(-1)) \\ &= M^\mathcal{T}(w^1(0)) * \overline{M^\mathcal{T}(w^2(0))} * \overline{M^\mathcal{T}(w^1(-1))} * M^\mathcal{T}(w^2(-1)) \\ &= M^\mathcal{T}(w^1(0)) * \overline{M^\mathcal{T}(w^2(0) + w^1(-1))} * M^\mathcal{T}(w^2(-1)) \\ &= M^\mathcal{T}(w^1(0)) * \left( \sum_{w'} u_{w,w'} x^{\tilde{B}v + \text{ind } C_q v} M^\mathcal{T}(w') \right) * M^\mathcal{T}(w^2(-1)), \end{aligned}$$

where we write  $w = w^2(0) + w^1(-1)$ ,  $w' = w - C_q v$ ,  $u_{w,w'} = u_{(0,w),(v,w)}$ . The monomial  $x^{\tilde{B}v + \text{ind } C_q v}$  is contained in  $\mathbb{Z}P[q^{\pm \frac{1}{2}}]$  and quasi-commutes with the other factors. Therefore, up to specified invertible elements in  $\mathbb{Z}[q^{\pm \frac{1}{2}}]$ , the above result becomes

$$\begin{aligned} & \sum_{w'} u_{w,w'} x^{\tilde{B}v + \text{ind } C_q v} M^\mathcal{T}(w^1(0)) * M^\mathcal{T}(w'(0)) * M^\mathcal{T}(w'(-1)) \\ & \quad * M^\mathcal{T}(w^2(-1)) \\ &= \sum_{w'} u_{w,w'} x^{\tilde{B}v + \text{ind } C_q v} M^\mathcal{T}(w^1(0) + w'(0) + w'(-1) + w^2(-1)). \end{aligned}$$

□

**Remark 7.1.11.** (1) *The above proof shows that the twisted product  $*$  of the standard basis elements is determined by Proposition 5.3.6 and 7.1.8.*

(2) *Notice that the map  $\widehat{\Pi\text{cor}}_{\chi_{q,t}}^{\leq 0}$  is algebraic when  $\tilde{Q}$  is of  $z$ -pattern. So Proposition 9.1.2 below provides another proof to Proposition 7.1.10.*

**7.2. Dual canonical basis elements.** For any  $l$ -dominant pair  $(v, w)$ , define  $L^\mathcal{T}_w(v) = \widehat{\Pi\text{cor}}_{\chi_{q,t}}^{\leq 0}(L_w(v))$  and denote it by  $L^\mathcal{T}(w)$  when  $v = 0$ . Then  $L^\mathcal{T}(w)$  is given by

$$L^\mathcal{T}(w) = \sum_v a_{v,0;w}(q^{\frac{\delta}{2}}) x^{\text{ind}(w) + \tilde{B}v},$$

where the Laurent polynomials  $a_{v,0;w}(t) = \sum_{d \in \mathbb{Z}} a_{v,0;w}^d t^d$  are given by 44. We shall see all the quantum cluster monomials essentially take this form.

**Lemma 7.2.1.** *We have*

$$L^\mathcal{T}_w(v) = L^\mathcal{T}(w - C_q v) x^{\tilde{B}v + \text{ind } C_q v}.$$

*Proof.* 1) Similar to (73), we compute

$$\begin{aligned} L^\mathcal{T}_w(v) &= \sum_{v'} \widehat{\Pi\text{cor}}(\langle L_w(v), \pi_w(v') \rangle) x^{\text{ind}(w)} x^{\tilde{B}v'} \\ &= \sum_{v'-v} \widehat{\Pi\text{cor}}(\langle L_{w-C_q v}(0), \pi_{w-C_q v}(v' - v) \rangle) x^{\text{ind}(w-C_q v)} x^{\tilde{B}(v'-v)} x^{\tilde{B}v + \text{ind}(w) - \text{ind}(w-C_q v)} \\ &= L^\mathcal{T}(w - C_q v) x^{\tilde{B}v + \text{ind } C_q v}. \end{aligned}$$

□

We already know that  $M_w(0) = \sum_v Z_{(0,w)(v,w)} L_w(v)$ . If we apply  $\widehat{\Pi\text{cor}} \widehat{\chi_{q,t}}^{\leq 0}$  to it, we obtain

$$M^\mathcal{T}(w) = \sum_v Z_{(0,w)(v,w)} L^\mathcal{T}(w - C_q v) x^{\tilde{B}v + \text{ind } C_q v}.$$

**7.3. Generic basis elements.** The generic basis elements  $\mathbb{L}^\mathcal{T}(w)$  are defined to be

$$(75) \quad \mathbb{L}^\mathcal{T}(w) = \sum_{v:(v,w) \text{ is } l\text{-dominant}} r_{w,w'} x^{\tilde{B}v + \text{ind } C_q v} L^\mathcal{T}(w - C_q v),$$

where  $w' = w - C_q v$ , and the integer  $r_{w,w'}$  is given in Definition 6.2.1. We have

$$\begin{aligned} \mathbb{L}^\mathcal{T}(w) &= \sum_v r_{w,w-C_q v} L^\mathcal{T}(w - C_q v) x^{\tilde{B}v + \text{ind } C_q v} \\ &= \sum_{v,v'} r_{w,w-C_q v} a_{v',0;w-C_q v}(q^{\frac{\delta}{2}}) x^{\text{ind}(w-C_q v) + \tilde{B}v'} x^{\tilde{B}v + \text{ind } C_q v} \\ &= \sum_{v,v'} r_{w,w-C_q v} a_{v'+v,v;w}(q^{\frac{\delta}{2}}) x^{\text{ind}(w)} x^{\tilde{B}(v+v')} \\ &= \sum_{v+v'=v''} b_{v''}(q^{\frac{\delta}{2}}) x^{\text{ind}(w)} x^{\tilde{B}v''}, \end{aligned}$$

where the coefficient  $b_{v''} = \sum_{(v,w) \text{ is } l\text{-dominant}} r_{w,w-C_q v} \cdot a_{v'',v;w}$ . Notice that  $b_{v''}$  is independent of the frozen pattern. Therefore, by taking the ice quiver  $\tilde{Q}$  to be of  $z$ -pattern, we obtain that  $b_{v''}$  equals the normalized Poincaré polynomial of the corresponding generic submodule Grassmannian.

## 8. BASES OF QUANTUM CLUSTER ALGEBRAS

We keep the assumptions on the ice quiver  $\tilde{Q}$  and the vectors  $v, w$  as in Section 7.

**8.1. Generic basis.** The results in Section 6 can be adapted to the generic basis elements  $\mathbb{L}^{\mathcal{T}}(w)$  in  $\mathcal{T}$ . We have the following theorem.

**Theorem 8.1.1.** 1) *The generic basis elements have the following expansion*

$$(76) \quad \mathbb{L}^{\mathcal{T}}(w) = \sum_v P_{q^{\frac{1}{2}\delta}}(\mathrm{Gr}_v(\sigma W)) q^{-\frac{1}{2} \dim \mathrm{Gr}_v(\sigma W)} x^{\mathrm{ind} w + \tilde{B}v}.$$

2) *We have a factorization*

$$(77) \quad \mathbb{L}^{\mathcal{T}}(w) = \mathbb{L}^{\mathcal{T}}(\phi w) \cdot \mathbb{L}^{\mathcal{T}}(f w).$$

3) *We have  $\mathbb{L}^{\mathcal{T}}(w) = L^{\mathcal{T}}(w)$  if a generic module of dimension  $\sigma w$  is rigid.*

In the presentable cluster category introduced by [Pla10a], cf. [Qin10, 2.3], Section 2.3, each  $w \in \mathcal{J}$  is associated with a coefficient-free generic object  $O(w)$ . We define the associated *generic quantum cluster characters* to be

$$(78) \quad \begin{aligned} \mathbb{L}^{\mathcal{A}}(w) &= \mathbb{L}^{\mathcal{T}}(w) \cdot x^{\mathrm{ind} O(w) - \mathrm{ind} w} \\ &= \sum_v P_{q^{\frac{1}{2}\delta}}(\mathrm{Gr}_v(\sigma W)) q^{-\frac{1}{2} \dim \mathrm{Gr}_v(\sigma W)} x^{\mathrm{ind} O(w) + \tilde{B}v}. \end{aligned}$$

**Remark 8.1.2.** We can denote  $\mathbb{L}^{\mathcal{A}}(w)$  by  $X_{O(w)}$ . This definition naturally generalizes the quantum cluster character formula in [Qin10, Definition 1.2.1] to generic objects.

It is not clear if one can extend this quantum character to arbitrary objects of the presentable cluster category in a reasonable way, for example, such that its image is still contained in  $\mathcal{A}^q$ . Some results for Dynkin quivers are discussed in [Din10].

**Proposition 8.1.3.** For any  $w \in \mathcal{J}$ , we have  $\mathrm{pr}_n \mathrm{ind} O(w) = \mathrm{pr}_n \mathrm{ind} w$ . When the ice quiver  $\tilde{Q}$  is the level 1 ice quiver with  $z$ -pattern, we have  $\mathrm{ind} O(w) = \mathrm{ind} w$ .

*Proof.* Because  $\mathrm{pr}_n \mathrm{ind} O(w)$  and  $\mathrm{pr}_n \mathrm{ind} w$  do not depend on the coefficients  $\tilde{Q} - Q$ , it suffices to prove the second statement.

First, the index  $\mathrm{ind} w$  is linear in  $w$ .

Second, the index  $\mathrm{ind} O(w)$  is linear with respects to the components in the canonical decomposition of the generic objects  $O(w)$ , which are either generic modules or the object  $P_i[1]$ ,  $i \in I$ .

Therefore, it suffices to study the indices of modules. Let  $O(w) = M$  be any  $Q^{op}$ -module with the minimal injective resolution

$$(79) \quad 0 \rightarrow M \xrightarrow{f} I^{w(-1)} \xrightarrow{g} I^{w(0)} \rightarrow 0.$$

Let  $B$  denote the Jacobi algebra of  $(\tilde{Q}, \tilde{W})$ . View the resolution as a short exact sequence in the category of  $B^{op}$ -modules. Denote the simple  $B^{op}$ -modules by  $S_j$ ,  $1 \leq j \leq 2n$ . If we apply  $\mathrm{Hom}_{B^{op}-\mathrm{mod}}(S_j, \_)$  to the above short exact sequence, we obtain a long exact sequence

$$\begin{aligned} \dots \rightarrow \mathrm{Hom}_{B^{op}-\mathrm{mod}}(S_j, M) &\xrightarrow{f_0} \mathrm{Hom}_{B^{op}-\mathrm{mod}}(S_j, I^{w(-1)}) \rightarrow \mathrm{Hom}_{B^{op}-\mathrm{mod}}(S_j, I^{w(0)}) \\ &\xrightarrow{w_0} \mathrm{Ext}_{B^{op}-\mathrm{mod}}^1(S_j, M) \xrightarrow{g_1} \mathrm{Ext}_{B^{op}-\mathrm{mod}}^1(S_j, I^{w(-1)}) \rightarrow \mathrm{Ext}_{B^{op}-\mathrm{mod}}^1(S_j, I^{w(0)}) \rightarrow \dots \end{aligned}$$

Then  $(\text{ind } M)_j = -\dim \text{Hom}_{B^{op}\text{-mod}}(S_j, M) + \dim \text{Ext}_{B^{op}\text{-mod}}^1(S_j, M)$ , cf. [Pal08].

Because  $M$  is supported on the principal part  $Q$ ,  $\dim \text{Hom}_{B^{op}\text{-mod}}(S_j, M)$  vanishes unless  $1 \leq j \leq n$ . The fact that (79) is the minimal injective resolution of  $Q^{op}$ -modules implies  $f_0$  is an isomorphism. It follows that  $w_0$  is injective.

When the quiver is of  $z$ -pattern, we have  $\text{ind } I_i = -e_i + e_{i+n}$ , cf. Lemma 7.1.1. Therefore  $\text{Ext}_{B^{op}\text{-mod}}^1(S_j, I^{w(-1)})$  equals  $w_{j-n}(-1)$  if  $j > n$  and vanishes elsewhere. Furthermore, the fact that (79) is the minimal injective resolution of the  $Q^{op}$ -module  $M$  implies that for any  $i \in I$ , either  $\dim \text{Ext}_{B^{op}\text{-mod}}^1(S_{i+n}, I^{w(-1)}) = w_i(-1)$  or  $\dim \text{Ext}_{B^{op}\text{-mod}}^1(S_{i+n}, I^{w(0)}) = w_i(0)$  is zero. Therefore,  $g_1$  is surjective.

Therefore, we have

$$\begin{aligned} (\text{ind } M)_j &= -\dim \text{Hom}_{B^{op}\text{-mod}}(S_j, I^{w(-1)}) + \dim \text{Hom}_{B^{op}\text{-mod}}(S_j, I^{w(0)}) \\ &\quad + \dim \text{Ext}_{B^{op}\text{-mod}}^1(S_j, I^{w(0)}) \\ &= \begin{cases} -w_j(-1) + w_j(0) & \text{if } j \leq n \\ w_j(0) + w_{j-n}(-1) & \text{if } j > n, \end{cases} \\ &= (\text{ind } w)_j. \end{aligned}$$

□

**Remark 8.1.4** (Failure 3). For general  $\tilde{Q}$ , we have

$$(80) \quad \text{ind } w \neq \text{ind } O(w).$$

**Lemma 8.1.5.** The elements in the coefficient ring  $\mathbb{Z}P[q^{\pm \frac{1}{2}}]$  quasi-commute with  $M^\mathcal{T}(w)$ ,  $L^\mathcal{T}(w)$ ,  $\mathbb{L}^\mathcal{T}(w)$ ,  $w \in \mathbb{N}^{I \times \{-1, 0\}}$ .

*Proof.* It follows from the fact that  $x_i$ ,  $n < i \leq N$ , commutes with  $x^{\tilde{B}v}$  for any  $v \in \mathbb{N}^n$ . □

**Proposition 8.1.6.** The set  $\{\mathbb{L}^\mathcal{T}(w), w \in \mathcal{J}\}$  is a  $\mathbb{Z}P[q^{\pm \frac{1}{2}}]$ -basis of the vector space  $\mathcal{A}_{sub}^q$ .

*Proof.* Notice that the matrix  $(r_{w, w'})$  is upper unitriangular. Furthermore, each element  $\mathbb{L}^\mathcal{T}(w)$  divides into the composition of the coefficient free part  $\mathbb{L}^\mathcal{T}(\phi w)$  and the pure coefficient part  $\mathbb{L}^\mathcal{T}(f w) \in \mathbb{Z}P[q^{\pm \frac{1}{2}}]$ . Therefore,  $\mathcal{A}_{sub}^q$  is spanned by  $\{\mathbb{L}^\mathcal{T}(w), w \in \mathcal{J}\}$  over  $\mathbb{Z}P[q^{\pm \frac{1}{2}}]$ .

Take any vectors  $w, w'$  in  $\mathcal{J}$ , such that  $w \neq w'$ . By proposition 8.1.3, the vector  $\text{pr}_n \text{ind}^z w$  is the index of a generic object in the cluster category of  $Q$ . Furthermore, we have  $\text{pr}_n \text{ind } w \neq \text{pr}_n \text{ind } w'$ , cf. the proof of Proposition 8.1.3 or [Pla10b]. Because  $\text{ind } w$  is the leading term of  $\mathbb{L}^\mathcal{T}(w)$  and  $\tilde{B}$  is of full rank, the set  $\{\mathbb{L}^\mathcal{T}(w), w \in \mathcal{J}\}$  is  $\mathbb{Z}P[q^{\pm \frac{1}{2}}]$ -linearly independent. □

**Proposition 8.1.7.**  $\mathcal{A}_{sub}^q$  equals  $\mathcal{A}^q$ .

*Proof.*  $\mathcal{A}_{sub}^q$  is a subspace of  $\mathcal{A}^q$  because the latter contains  $\{M^\mathcal{T}(w)\}$ . If we take any two elements  $\mathbb{L}^\mathcal{T}(\phi w^1) * f_1, \mathbb{L}^\mathcal{T}(\phi w^2) * f_2$ , where  $f_1, f_2 \in \mathbb{Z}P[q^{\pm \frac{1}{2}}]$ , it follows from Proposition 7.1.10 that their twisted product still belongs

to  $\mathcal{A}_{sub}^q$ . Therefore,  $\mathcal{A}_{sub}^q$  is a subalgebra of  $\mathcal{A}^q$  with respect to the twisted product. But it contains all the quantum cluster variables. Therefore, the two algebras must agree.  $\square$

**Theorem 8.1.8.**  $\{\mathbb{L}^{\mathcal{A}}(w), w \in \mathcal{J}\}$  is a  $\mathbb{Z}P[q^{\pm \frac{1}{2}}]$ -basis of  $\mathcal{A}^q$ . Furthermore, it contains all the quantum cluster monomials.

*Proof.* The statement follows from Proposition 8.1.6 and 8.1.7.  $\square$

**8.2. Dual canonical basis and dual PBW basis.** Similar to the treatment of generic quantum cluster characters, for each  $w \in \mathcal{J}$ , we normalize the quantum torus elements  $M^{\mathcal{T}}(w)$  and  $L^{\mathcal{T}}(w)$  by defining

$$(81) \quad M^{\mathcal{A}}(w) = M^{\mathcal{T}}(w) \cdot x^{\text{ind } O(w) - \text{ind } w},$$

$$(82) \quad L^{\mathcal{A}}(w) = L^{\mathcal{T}}(w) \cdot x^{\text{ind } O(w) - \text{ind } w}.$$

$$(83)$$

**Theorem 8.2.1.** The sets  $\{M^{\mathcal{A}}(w), w \in \mathcal{J}\}$ ,  $\{L^{\mathcal{A}}(w), w \in \mathcal{J}\}$  are  $\mathbb{Z}P[q^{\pm \frac{1}{2}}]$ -bases of  $\mathcal{A}^q$ . They are called the dual PBW basis and the dual canonical basis of the quantum cluster algebra  $\mathcal{A}^q$  respectively. Furthermore, all the quantum cluster monomials are contained in  $\{L^{\mathcal{A}}(w), w \in \mathcal{J}\}$ .

*Proof.* It suffices to show that  $\{M^{\mathcal{T}}(w), w \in \mathcal{J}\}$ ,  $\{L^{\mathcal{T}}(w), w \in \mathcal{J}\}$  are  $\mathbb{Z}P[q^{\pm \frac{1}{2}}]$ -bases of  $\mathcal{A}^q$ . Applying the truncated character map to (46), we obtain

$$(84) \quad M^{\mathcal{T}}(w) = L^{\mathcal{T}}(w) + \sum_{(v,w) \text{ is } l\text{-dominant}} Z_{w,w'}(q^{\frac{\delta}{2}}) x^{\tilde{B}v + \text{ind } C_q v} L^{\mathcal{T}}(w'),$$

where  $w' = w - C_q v$  and  $Z_{w,w'}(t) = Z_{0,v;w}(t) \in t^{-1}\mathbb{Z}[t^{-1}]$ . Denote the matrix of the coefficients  $Z_{w,w'}(q^{\frac{\delta}{2}}) x^{\tilde{B}v + \text{ind } C_q v}$  by  $\tilde{Z}(q^{\frac{\delta}{2}})$ .

Similarly, denote the matrix of the coefficients  $r_{w,w'} x^{\tilde{B}v + \text{ind } C_q v}$  in (75) by  $R$  and its inverse by  $R^{-1}$ .

Define the  $\mathcal{J} \times \mathcal{J}$  matrix  ${}^{\phi}R^{-1}$  such that its entry in position  $(w, w')$  is

$$({}^{\phi}R^{-1})_{w,w'} = \sum_{w'' \in \mathbb{N}^I \times \{-1, 0\}; {}^{\phi}w'' = w'} R_{w,w''}^{-1} x^{\text{ind}(f w'')}.$$

Denote the product  $\tilde{Z}(q^{\frac{\delta}{2}})R^{-1}$  by  $S(q^{\frac{\delta}{2}})$ . Similarly, the  $\mathcal{J} \times \mathcal{J}$  matrix  ${}^{\phi}S(q^{\frac{\delta}{2}})$  is defined such that its entry in position  $(w, w')$  is

$$({}^{\phi}S(q^{\frac{\delta}{2}}))_{w,w'} = \sum_{w'' \in \mathbb{N}^I \times \{-1, 0\}; {}^{\phi}w'' = w'} S(q^{\frac{\delta}{2}})_{w,w''} x^{\text{ind}(f w'')}.$$

The matrix transition between  $\{\mathbb{L}^{\mathcal{T}}(w), w \in \mathcal{J}\}$  and  $\{L^{\mathcal{T}}(w), w \in \mathcal{J}\}$  is given by the matrix  ${}^{\phi}R^{-1}$ . The matrix transition between  $\{\mathbb{L}^{\mathcal{T}}(w), w \in \mathcal{J}\}$  and  $\{M^{\mathcal{T}}(w), w \in \mathcal{J}\}$  is given by the matrix  ${}^{\phi}S(q^{\frac{\delta}{2}})$ .

With respect to the dominance order, the matrices  $R^{-1}$  and  $S$  are upper unitriangular. It follows that  ${}^{\phi}R^{-1}$  and  ${}^{\phi}S$  are upper triangular. Notice that, if for  $w, w'$  in  $\mathbb{N}^I \times \{-1, 0\}$ , we have  $w' \leq w$  with respect to the dominance order, then  $w'$  must be equal to  $w$ . It follows that  ${}^{\phi}R^{-1}$  and  ${}^{\phi}S$  are upper unitriangular. Therefore, we obtain the statements from Theorem 8.1.8.  $\square$

**Remark 8.2.2.** *The transition matrices vary little when the choice of the coefficients and quantization change:*

1) *The matrix  $Z$  can be solved recursively by a combinatorial algorithm, cf. [Lus90, 7.10] or [Nak04, Section 8].  $Z$  only depends on  $Q$ . Also, the geometrically defined integer matrix  $r$  only depends on  $Q$ . The set  $\{w' : \phi w' = w''\}$  for given  $w''$  can be calculated easily.*

2) *We have  $\tilde{Z}(q^{\frac{\delta}{2}}) = Z(q^{\frac{\delta}{2}})$  and  $R = r$  if the quiver is of  $z$ -pattern.*

**8.3. Structure constants.** In [Kim12], Kimura studied the factorization of the dual canonical basis  $\mathbf{B}^{\text{up}}$  of a certain quantum unipotent subgroup. Because in [KQ12], the authors identified  $L(w)$  with elements in  $\mathbf{B}^{\text{up}}$ , we can translate his result into our setting.

**Theorem 8.3.1** ([Kim12, Theorem 6.21]). *Up to  $t$ -powers, we have the factorization of simples*

$$(85) \quad L(w) = L(\phi w) \cdot L(f w).$$

For any  $w^1, w^2, w^3$  in  $\mathcal{J}$ , define an element in  $\mathbb{Z}P[q^{\pm \frac{1}{2}}]$

$$(86) \quad \phi b_{w^1, w^2; \tilde{B}}^{w^3} = x^{\text{ind } O(w^1) + \text{ind } O(w^2) - \text{ind } O(w^3) - \text{ind}(w^1 + w^2 - w^3)} \sum_{w: \phi w = w^3} b_{w^1, w^2}^w x^{\text{ind } f w} x^{\tilde{B}v + \text{ind } C_q v}$$

where  $v$  is determined by

$$(87) \quad w = w^1 + w^2 - C_q v.$$

**Theorem 8.3.2** (Positive basis). *The structure constants of the dual canonical basis  $\{L^\mathcal{T}(w), w \in \mathcal{J}\}$  are given by, for any  $w^1, w^2$  in  $\mathcal{J}$ ,*

$$L^\mathcal{A}(w^1) * L^\mathcal{A}(w^2) = q^{\frac{1}{2}\Lambda(\text{ind}(w^1), \text{ind}(w^2)) - \frac{\delta}{2}\Lambda^z(\text{ind}^z(w^1), \text{ind}^z(w^2))} \cdot \sum_{w^3 \in \mathcal{J}} \phi b_{w^1, w^2; \tilde{B}}^{w^3}(q^{\frac{\delta}{2}}) L^\mathcal{A}(w^3).$$

*In particular, they are contained in  $\mathbb{N}[q^{\pm \frac{1}{2}}][x_{n+1}^\pm, \dots, x_m^\pm]$ .*

*Proof.* First consider the quantum cluster algebra  $\mathcal{A}^q$  associated with  $(\tilde{B}^z, \Lambda^z)$ . Then we have  $\widehat{\Pi} \chi_{q,t} = \chi_{q,t} \widehat{\Pi}$ . The truncated character  $\chi_{q,t}^{\leq 0}$  is algebraic from  $R_t$  to  $D(\mathcal{T})$ . Therefore, the  $\mathbb{Z}[t^\pm]$ -linear independent set  $\{L^\mathcal{T}(w)\}$  in  $\mathcal{T}$  is closed under the twisted product  $*$  and the corresponding positive structure constants are  $b_{w^1, w^2}^w$  as given by (47). Using Theorem 8.3.1, we obtain the structure constants of the dual canonical basis  $\{L^\mathcal{T}(w), w \in \mathcal{J}\}$ .

Finally, the correction technique in Proposition 9.1.2 implies the above result for any acyclic quantum cluster algebra.  $\square$

We obtain that, whatever the coefficient pattern  $\tilde{Q} - Q$  and the quantization  $\Lambda$  we choose, the quantum cluster algebras containing acyclic seeds are *strongly positive*: they admit a basis in which the structure constants are positive.

The following important consequence of the deformed monoidal categorification is an easy generalization of the first main result of the author's joint work with Kimura [KQ12] for arbitrary choice of coefficients and quantization.

**Corollary 8.3.3.** (*Quantum positivity [KQ12]*) Any quantum cluster monomial  $m$  can be written as a Laurent polynomial of the quantum cluster variables  $x_i$ ,  $1 \leq i \leq n$ , in any given seed with coefficients in  $\mathbb{N}[q^{\pm \frac{\delta}{2}}, x_{n+1}^{\pm}, \dots, x_m^{\pm}]$ .

*Proof.* By the quantum Laurent phenomenon, we have

$$m = \frac{\sum_{m_*=(m_i)} c_{m_*} \prod_{1 \leq i \leq n} x_i^{m_i}}{\prod_i x_i^{d_i}},$$

where  $m_* = (m_i)_{i \in I}$ ,  $d_* = (d_i)_{i \in I}$  are sequences of nonnegative integers and the coefficients  $c_{m_*}$  are contained in  $\mathbb{Z}P[q^{\pm \frac{1}{2}}]$ . Notice that we use the usual product  $\cdot$  in this expression.

The quantum cluster monomial  $m$  equals  $L^{\mathcal{A}}(w)$  for some  $w$ . Also, the quantum  $X$ -variable  $x_i$ ,  $1 \leq i \leq m$ , equals  $L^{\mathcal{A}}(w_i)$  for some  $w_i$ . We can rewrite the above equation as

$$\begin{aligned} \sum_{m_*=(m_i)} c_{m_*} L^{\mathcal{A}}\left(\sum_i m_i w_i\right) &= \prod_i L^{\mathcal{A}}(w_i)^{d_i} \cdot L^{\mathcal{A}}(w) \\ &= q^{-\frac{1}{2}\Lambda(\text{ind}(\sum_i d_i w_i), \text{ind}(w))} L^{\mathcal{A}}\left(\sum_i d_i w_i\right) * L^{\mathcal{A}}(w). \end{aligned}$$

The statement follows from Theorem (8.3.2).  $\square$

The almost simple pseudo-modules  $\mathbb{L}(w)$  introduced in Section 6 form a  $\mathbb{Z}[q^{\pm \frac{1}{2}}]$ -basis of  $\mathcal{R}_t$ . Notice that, when the ice quiver is of  $z$ -pattern, we have  $\widehat{\Pi}\text{cor}_{\chi_{q,t} \leq 0} \mathbb{L}(w) = \mathbb{L}^{\mathcal{T}}(w) = \mathbb{L}^{\mathcal{A}}(w)$ . Denote the structure constants of this basis by  $c_{w^1, w^2}^w(t)$ : for any  $w^1, w^2$  in  $\mathbb{N}^{I \times \{-1, 0\}}$ , we have

$$(88) \quad \mathbb{L}(w^1) \otimes \mathbb{L}(w^2) = \sum_w c_{w^1, w^2}^w \mathbb{L}(w).$$

Because the matrix  $(r_{w, w'})$  and the matrix of the structure constants of  $\{\mathbb{L}^{\mathcal{T}}(w)\}$  are upper unitriangular, the matrix  $(c_{w^1, w^2}^w)$  is upper unitriangular as well. Therefore, for every nonzero term in the sum, we always have  $w = w^1 + w^2 - C_q v$  for some  $v$ .

For any  $w^1, w^2, w^3$  in  $\mathcal{J}$ , define an element in  $\mathbb{Z}P[q^{\pm \frac{1}{2}}]$

$$(89) \quad \phi_{c_{w^1, w^2; \tilde{B}}}^{w^3} = x^{\text{ind } O(w^1) + \text{ind } O(w^2) - \text{ind } O(w^3) - \text{ind}(w^1 + w^2 - w^3)} \sum_{w: \phi w = w^3} c_{w^1, w^2}^w x^{\text{ind } f w} x^{\tilde{B}v + \text{ind } C_q v}$$

where  $v$  is determined by  $w = w^1 + w^2 - C_q v$ .

**Theorem 8.3.4.** The structure constants of the generic basis  $\{\mathbb{L}^{\mathcal{T}}(w), w \in \mathcal{J}\}$  are given by, for any  $w^1, w^2$  in  $\mathcal{J}$ ,

$$\begin{aligned} \mathbb{L}^{\mathcal{A}}(w^1) * \mathbb{L}^{\mathcal{A}}(w^2) &= q^{\frac{1}{2}\Lambda(\text{ind}(w^1), \text{ind}(w^2)) - \frac{\delta}{2}\Lambda^z(\text{ind}^z(w^1), \text{ind}^z(w^2))} \\ &\quad \cdot \sum_{w^3 \in \mathcal{J}} \phi_{c_{w^1, w^2; \tilde{B}}}^{w^3} (q^{\frac{\delta}{2}}) \mathbb{L}^{\mathcal{A}}(w^3). \end{aligned}$$

*Proof.* This theorem is a consequence of (77) and Proposition 9.1.2. The proof is similar to that of Theorem 8.3.2.  $\square$

## 9. CHANGE OF COEFFICIENTS AND QUANTIZATIONS

**9.1. Correction technique.** Given any (quantum) cluster algebra, if some basis of it is known, we want to ask what happens if we change the coefficients and the quantization. As we have seen in Theorem 8.1.8 Theorem 8.2.1 and Remark 8.2.2, the bases and their transition matrices vary little the  $\tilde{Q} - Q$  and  $\Lambda$  change. We shall show that this phenomenon is true in general, by generalizing the correction factors in the previous failures (70) (73) (80).

Let  $n, m^{(1)}, m^{(2)}$  be three integers, such that  $0 < n \leq m^{(1)}, m^{(2)}$ . For  $i = 1, 2$ , let  $\tilde{B}^{(i)}$  be a rank  $m^{(i)} \times n$  matrix. Let  $\Lambda^{(i)}$  be a rank  $m^{(i)} \times m^{(i)}$  skew-symmetric integer matrix.

**Definition 9.1.1** (weakly compatible pair). *The pair  $(\tilde{B}^{(i)}, \Lambda^{(i)})$  is called weakly compatible if there exists an  $n \times n$  integer matrix  $D^{(i)}$  such that we have*

$$(90) \quad \Lambda^{(i)}(-\tilde{B}^{(i)}) = \begin{bmatrix} D^{(i)} \\ 0 \end{bmatrix}.$$

Assume the matrices  $\tilde{B}^{(i)}$  have the common principal part  $B$ , the pairs  $(\tilde{B}^{(i)}, \Lambda^{(i)})$  are weakly compatible, and  $D^{(2)} = \delta D^{(1)}$  for some integer  $\delta$ . A priori,  $\text{rank } D^{(1)}$  is no less than  $\text{rank } D^{(2)}$ .

Define the associated quantum tori

$$\mathcal{T}^{(i)} = \mathcal{T}^{(i)}(\Lambda^{(i)}) = \mathbb{Z}[q^{\pm \frac{1}{2}}][x_1^{\pm}, \dots, x_{m^{(i)}}^{\pm}]$$

such that twisted product is determined by  $\Lambda^{(i)}$ .

Let  $\max(m^{(1)}, m^{(2)})$  be denoted by  $m$ . As in (68), enlarge  $\tilde{B}^{(1)}, \tilde{B}^{(2)}$  into  $m \times n$  matrices and  $\Lambda^{(2)}$  into  $m \times m$  matrix by adding zero entries, and define the associated enlarged quantum torus

$$\mathcal{T} = \mathcal{T}(\Lambda^{(2)}) = \mathbb{Z}[q^{\pm \frac{1}{2}}][x_1^{\pm}, \dots, x_m^{\pm}],$$

such that its twisted product is determined by the enlarged matrix  $\Lambda^{(2)}$ .

For each  $i$ , and for any integer  $s \geq 3$ , integer  $1 \leq j \leq s$ , vector  $g^{(i)}_j \in \mathbb{Z}^{n_i}$ , and polynomial

$$F_j(t; y_1, \dots, y_n) = \sum_{v \in \mathbb{N}^n} b_j(t; v) y^v \in \mathbb{Z}[t^{\pm}][y_1, \dots, y_n],$$

where  $t, y_1, \dots, y_n$  are indeterminates,  $b_j(t; v) \in \mathbb{Z}[t^{\pm}]$ , we define the following element in the quantum torus  $\mathcal{T}^{(1)}$ :

$$\begin{aligned} M_j^{(1)} &= x_j^{g_j^{(1)}} F_j(q^{\frac{1}{2}}; x^{\tilde{B}^{(1)} e_1}, \dots, x^{\tilde{B}^{(1)} e_n}) \\ &= x_j^{g_j^{(1)}} \sum_{v \in \mathbb{N}^n} b_j(q^{\frac{1}{2}}; v) x^{\tilde{B}^{(1)} v}, \end{aligned}$$

where  $e_k$  is the  $k$ -th unite vector in  $\mathbb{Z}^n$ , and a similar element in  $\mathcal{T}^{(2)}$ :

$$\begin{aligned} M_j^{(2)} &= x_j^{g_j^{(2)}} F_j(q^{\frac{\delta}{2}}; x^{\tilde{B}^{(2)} e_1}, \dots, x^{\tilde{B}^{(2)} e_n}) \\ &= x_j^{g_j^{(2)}} \sum_{v \in \mathbb{N}^n} b_j(q^{\frac{\delta}{2}}; v) x^{\tilde{B}^{(2)} v}. \end{aligned}$$

Assume the first  $n$  coordinates of  $g_j^{(1)}$  and  $g_j^{(2)}$  are equal, *i.e.* we have  $\text{pr}_n g_j^{(1)} = \text{pr}_n g_j^{(2)} = g_j$  for some vector  $g_j \in \mathbb{Z}^n$ .

**Proposition 9.1.2.** *Assume that  $\tilde{B}^{(1)}$  is of full rank and  $b_j(t; 0)$  does not vanish for any  $3 \leq j \leq s$ . If the following equation holds in  $\mathcal{T}^{(1)}$ :*

$$(91) \quad q^{-\frac{1}{2}\Lambda^{(1)}(g_1^{(1)}, g_2^{(1)})} M_1^{(1)} * M_2^{(1)} = \sum_{3 \leq j \leq s} c_j^{(1)}(q^{\frac{1}{2}}) M_j^{(1)}$$

for some coefficients  $c_j^{(1)}(t) \in \mathbb{Z}[t^\pm][x_{n+1}, \dots, x_{m(1)}^\pm]$ , then there exist unique vectors  $u_j \in \mathbb{N}^n$  such that  $g_j^{(1)} = g_1^{(1)} + g_2^{(1)} + \tilde{B}^{(1)} u_j$ , and we have

$$(92) \quad q^{-\frac{1}{2}\Lambda^{(2)}(g_1^{(2)}, g_2^{(2)})} M_1^{(2)} * M_2^{(2)} = \sum_{3 \leq j \leq s} c_j^{(2)}(q^{\frac{\delta}{2}}) M_j^{(2)}$$

such that the coefficients in  $\mathbb{Z}[t^\pm][x_{n+1}, \dots, x_{m(2)}^\pm]$  are given by

$$(93) \quad c_j^{(2)}(q^{\frac{\delta}{2}}) = c_j^{(1)}(q^{\frac{\delta}{2}}) x^{g_1^{(2)} + g_2^{(2)} + \tilde{B}^{(2)} u_j - g_j^{(2)}}.$$

*Proof.* Expanding LHS of (91), we obtain

$$\begin{aligned} \text{LHS} &= \sum_{v_1, v_2: v_1 + v_2 = v} q^{-\frac{1}{2}\Lambda^{(1)}(g_1^{(1)}, g_2^{(1)}) + \frac{1}{2}\Lambda^{(1)}(g_1^{(1)} + \tilde{B}^{(1)} v_1, g_2^{(1)} + \tilde{B}^{(1)} v_2)} \\ &\quad \cdot b_1(q^{\frac{1}{2}}; v_1) b_2(q^{\frac{1}{2}}; v_2) x^{g_1^{(1)} + g_2^{(1)} + \tilde{B}^{(1)} v} \\ &= \sum_{v_1, v_2: v_1 + v_2 = v} q^{\frac{1}{2}(-g_1^T D^{(1)} v_2 + g_2^T D^{(1)} v_1 + v_1^T B D^{(1)} v_2)} \\ &\quad \cdot b_1(q^{\frac{1}{2}}; v_1) b_2(q^{\frac{1}{2}}; v_2) x^{g_1^{(1)} + g_2^{(1)} + \tilde{B}^{(1)} v} \\ &= \sum_{v_1, v_2: v_1 + v_2 = v} q^{\frac{1}{2\delta}(-g_1^T D^{(2)} v_2 + g_2^T D^{(2)} v_1 + v_1^T B D^{(2)} v_2)} \\ &\quad \cdot b_1(q^{\frac{1}{2}}; v_1) b_2(q^{\frac{1}{2}}; v_2) x^{g_1^{(1)} + g_2^{(1)} + \tilde{B}^{(1)} v}. \end{aligned}$$

In (91), since  $b_j(t; 0)$  is nonzero, the monomial  $x^{g_j^{(1)}}$  in RHS must be killed by either another monomial  $x^{g_{j'}^{(1)} + \tilde{B}^{(1)} v_{j'}}$  in RHS or a monomial in LHS. In the former case, repeat this argument for  $x^{g_{j'}^{(1)}}$ . Then, after finite steps, we obtain that for some  $u_j \in \mathbb{N}^n$ ,  $x^{g_j^{(1)}}$  must equal the monomial  $x^{g_1^{(1)} + g_2^{(1)} + \tilde{B}^{(1)} u_j}$  in LHS. Therefore, we can rewrite RHS as

$$\text{RHS} = \sum_j c_j^{(1)}(q^{\frac{1}{2}}) \sum_{v_j} b_j(q^{\frac{1}{2}}; v_j) x^{g_1^{(1)} + g_2^{(1)} + \tilde{B}^{(1)}(u_j + v_j)}.$$

Viewed the both sides as usual Laurent polynomial and embed them into the enlarged quantum torus  $\mathcal{T}$ . We can rewrite them as

$$\begin{aligned} \text{LHS} &= \sum_{v_1, v_2: v_1 + v_2 = v} (q^{\frac{1}{2\delta}})^{-\Lambda^{(2)}(g_1^{(2)}, g_2^{(2)}) + \Lambda^{(2)}(g_1^{(2)} + \tilde{B}^{(2)}v_1, g_2^{(2)} + \tilde{B}^{(2)}v_2)} \\ &\quad \cdot b_1(q^{\frac{1}{2}}; v_1) b_2(q^{\frac{1}{2}}; v_2) x^{g_1^{(2)} + g_2^{(2)} + \tilde{B}^{(2)}v} \\ &\quad \cdot x^{g_1^{(1)} + g_2^{(1)} - g_1^{(2)} - g_2^{(2)}} x^{(\tilde{B}^{(1)} - \tilde{B}^{(2)})v}, \\ \text{RHS} &= \sum_{j, v_j: u_j + v_j = v} c_j^{(1)}(q^{\frac{1}{2}}) b_j(q^{\frac{1}{2}}; v_j) x^{g_1^{(2)} + g_2^{(2)} + \tilde{B}^{(2)}(u_j + v_j)} \\ &\quad \cdot x^{g_1^{(1)} + g_2^{(1)} - g_1^{(2)} - g_2^{(2)}} x^{(\tilde{B}^{(1)} - \tilde{B}^{(2)})(u_j + v_j)}. \end{aligned}$$

In the both sides, we replace the indeterminate  $q^{\frac{1}{2}}$  by  $q^{\frac{\delta}{2}}$  and delete the factor

$$x^{g_1^{(1)} + g_2^{(1)} - g_1^{(2)} - g_2^{(2)}} x^{(\tilde{B}^{(1)} - \tilde{B}^{(2)})v}$$

in each monomial. Then we still have  $\text{LHS} = \text{RHS}$ , which now become

$$\begin{aligned} \text{LHS} &= q^{-\frac{1}{2}\Lambda^{(2)}(g_1^{(2)}, g_2^{(2)})} M_1^{(2)} * M_2^{(2)} \\ \text{RHS} &= \sum_j \sum_{v_j} c_j^{(1)}(q^{\frac{\delta}{2}}) b_j(q^{\frac{\delta}{2}}; v_j) x^{g_1^{(2)} + g_2^{(2)} + \tilde{B}^{(2)}u_j - g_j^{(2)}} x^{g_j^{(2)} + \tilde{B}^{(2)}v_j} \\ &= \sum_j c_j^{(1)}(q^{\frac{\delta}{2}}) x^{g_1^{(2)} + g_2^{(2)} + \tilde{B}^{(2)}u_j - g_j^{(2)}} M_j. \end{aligned}$$

Thus, (92) is proved.  $\square$

**Remark 9.1.3.** The quotient between the  $q$ -powers on the left of (91) and (92) should be viewed as the generalization of the  $q$ -power correction factor in Failure 1 (70). The correction factor  $x^{g_1^{(2)} + g_2^{(2)} + \tilde{B}^{(2)}u_j - g_j^{(2)}}$  should be viewed as the generalization of the factor in Failure 2 (73). The difference  $g_j^{(2)} - g_j^{(1)}$  should be viewed as the generalization of the difference of the two sides of (80).

## 9.2. Change of (quantum) cluster algebras.

**Theorem 9.2.1.** For  $i = 1, 2$ , assume that the pair  $(\tilde{B}^{(i)}, \Lambda^{(i)})$ ,  $i = 1, 2$ , in Proposition 9.1.2 is either compatible or the matrix  $\Lambda^{(i)}$  is zero. Denote the corresponding quantum or classical cluster algebras by  $\mathcal{A}^{q(i)}$ .

1) If the (quantum) cluster variables of  $\mathcal{A}^{q(1)}$  take the form  $M_j^{(1)}$ , then the corresponding (quantum) cluster variables of  $\mathcal{A}^{q(2)}$  are given by the corresponding  $M_j^{(2)}$  such that  $g_j^{(2)}$  are the extended  $g$ -vectors.

2) For  $i = 1, 2$ , let  $\mathbf{B}^{(i)}$  be a  $\mathbb{Z}[q^{\pm \frac{1}{2}}][x_{n+1}^{\pm}, \dots, x_{m(i)}^{\pm}]$ -basis of  $\mathcal{A}^{q(i)}$  such that its elements take the form  $M^{(i)}_j$ . If the structure constants of  $\mathbf{B}^{(1)}$  are described by (91), then the structure constants of  $\mathbf{B}^{(2)}$  are described by the corresponding equation (92).

*Proof.* The first statement is obtained by taking (91) to be the quantum exchange relation.

The second statement is obtained by taking (91) to be the multiplication of basis elements.  $\square$

**Remark 9.2.2.** *By this theorem, in order to study quantum cluster variables and bases, it suffices to look at a special coefficient pattern with special quantizations, e.g. the principal coefficients with the canonical quantization.*

*For special coefficient patterns and special quantizations, by categorification techniques we can prove that the (quantum) cluster variables of  $\mathcal{A}^{q(1)}$  take the form  $M_j^{(1)}$ , cf. the commutative case by [CC06], [CK08], [DWZ10], [Pla10b], and the quantum case by [Qin10] when  $Q$  is acyclic and by [Nag10] in general. The first statement then implies that we have the (quantum) Laurent phenomenon and the existence of the (quantum)  $F$ -polynomials for general quantum cluster algebra  $\mathcal{A}^{q(2)}$ .*

*The first statement also implies Theorem 3.0.5 for general quantum cluster algebras.*

**9.3. Acyclic case.** Now we can apply the correction techniques to the results of Section 8. We simplify the proofs of previous results on bases of acyclic quantum cluster algebras, and we also present results about the bases of acyclic classical cluster algebras.

Let the compatible pairs  $(\tilde{B}^{(i)}, \Lambda^{(i)})$ ,  $i = 1, 2$ , be given as in Theorem 9.2.1. Further assume that  $B = B_Q$  for some acyclic quiver, and choose  $(\tilde{B}^{(1)}, \Lambda^{(1)})$  to be  $(\tilde{B}^z, \Lambda^z)$ . Then we can choose the  $\mathbb{Z}[q^{\pm\frac{1}{2}}]$ -linearly independent subset<sup>6</sup>  $\mathbf{B}^{(1)}$  of the elements of  $\mathcal{A}^{q(1)}$  to be the set  $\{\mathbb{L}^A(w)\}$ , the set  $\{M^A(w)\}$ , or the set  $\{L^A(w)\}$ . Notice that the elements of  $\mathbf{B}^{(1)}$  take the form  $M_j^{(1)}$  of Proposition 9.1.2. Define  $\mathbf{B}^{(2)}$  to be the subset  $\{\mathbb{L}^A(w), w \in \mathcal{J}\}$ ,  $\{M^A(w), w \in \mathcal{J}\}$ ,  $\{L^A(w), w \in \mathcal{J}\}$  in  $\mathcal{T}^{(2)}$  respectively if  $\Lambda^{(2)}$  is nonzero, or the corresponding classical limit if  $\Lambda^{(2)} = 0$ . Then the elements of  $\mathbf{B}^{(2)}$  take the form  $M_j^{(2)}$ .

**Theorem 9.3.1.** *If  $\tilde{B}^{(2)}$  is of full rank, then  $\mathbf{B}^{(2)}$  is a  $\mathbb{Z}P[q^{\pm\frac{1}{2}}]$ -basis of  $\mathcal{A}^{q(2)}$ . Furthermore, if  $\mathbf{B}^{(2)}$  is  $\{\mathbb{L}^A(w), w \in \mathcal{J}\}$  or  $\{L^A(w), w \in \mathcal{J}\}$ , then the structure constants of  $\mathbf{B}^{(2)}$  can be deduced from (92) which corresponds to the structure constants equation (91).*

*Proof.* By the proof of 8.1.6, the leading terms (terms with  $v = 0$ ) of the elements in  $\mathbf{B}^{(2)}$  are different. Because  $\tilde{B}^{(2)}$  is of full rank, in each  $M_j^{(2)}$ , the leading term can not be killed by the other terms. Therefore,  $\mathbf{B}^{(2)}$  is linearly independent.

Because  $\mathbf{B}^{(1)}$  is contained in  $\mathcal{A}^{q(1)}$ , by Proposition 9.1.2,  $\mathbf{B}^{(2)}$  is contained in  $\mathcal{A}^{q(2)}$ . Using Theorem 9.2.1 and (64) or (85), we obtain that  $\mathbf{B}^{(2)}$  is a  $\mathbb{Z}P[q^{\pm\frac{1}{2}}]$ -basis if it is either  $\{\mathbb{L}^A(w), w \in \mathcal{J}\}$  or  $\{L^A(w), w \in \mathcal{J}\}$ . The statement concerning its structure constants also follows from Proposition 9.1.2.

Using the upper unitriangular basis transition matrices studied before, we deduce that  $\{M^A(w), w \in \mathcal{J}\}$  is also a basis.  $\square$

<sup>6</sup>The subset  $\mathbf{B}^{(1)}$  is a basis of the subalgebra of  $\mathcal{A}^{q(1)}$  generated by the quantum cluster variables and the frozen variables  $x_{n+1}, \dots, x_{2n}$ .

**Remark 9.3.2.** If we take  $\Lambda^{(2)} = 0$ , then  $\mathcal{A}^{(2)}$  is a classical cluster algebra, which we denote by  $\mathcal{A}^{(2)}$ .

1) If we take  $\mathbf{B}^{(1)}$  to be the dual canonical basis, then we obtain that  $\mathbf{B}^{(2)}$  is the dual canonical basis of  $\mathcal{A}^{(2)}$  with positive structure constants. All cluster monomials are contained in  $\mathbf{B}^{(2)}$ .

2) If we take  $\mathbf{B}^{(1)}$  to be the generic basis and the dual PBW basis, then this theorem generalizes the results of the dual semicanonical basis and the dual PBW basis in [GLS11, 16.1] to the integral form and general coefficient type.

Recently, another proof of the generic basis for  $\mathcal{A}^{(2)}$  is obtained by [GLS12] [Pla11], which is based on cluster categories and a combinatorial result of [BFZ05].

## REFERENCES

- [Ami09] Claire Amiot, *Cluster categories for algebras of global dimension 2 and quivers with potential*, Annales de l’institut Fourier **59** (2009), no. 6, 2525–2590, [arXiv:0805.1035](#).
- [BBD82] Alexander A. Beilinson, Joseph Bernstein, and Pierre Deligne, *Analyse et topologie sur les espaces singuliers*, Astérisque, vol. 100, Soc. Math. France, 1982 (French).
- [BFZ05] Arkady Berenstein, Sergey Fomin, and Andrei Zelevinsky, *Cluster algebras. III. Upper bounds and double Bruhat cells*, Duke Math. J. **126** (2005), no. 1, 1–52.
- [BZ05] Arkady Berenstein and Andrei Zelevinsky, *Quantum cluster algebras*, Adv. Math. **195** (2005), no. 2, 405–455, [arXiv:math/0404446v2](#).
- [BZ11] ———, *Quantum cluster algebras*, lectures in Osaka, 2011.
- [BZ12] ———, *Triangular bases in quantum cluster algebras*, [arXiv:1206.3586](#).
- [CC06] Philippe Caldero and Frédéric Chapoton, *Cluster algebras as Hall algebras of quiver representations*, Comment. Math. Helv. **81** (2006), no. 3, 595–616, [arXiv:math.RT/0410187v2](#).
- [CK08] Philippe Caldero and Bernhard Keller, *From triangulated categories to cluster algebras*, Inv. Math. **172** (2008), 169–211.
- [CG97] N. Chriss and V. Ginzburg, *Representation theory and complex geometry*, Progress in Math., Birkhäuser, 1997.
- [CBJ05] William Crawley-Boevey and Bernt Tore Jensen, *A note on sub-bundles of vector bundles*, 2005, [arXiv:math/0505149v2](#).
- [DWZ08] Harm Derksen, Jerzy Weyman, and Andrei Zelevinsky, *Quivers with potentials and their representations I: Mutations*, Selecta Mathematica **14** (2008), 59–119, [arXiv:0704.0649v4](#).
- [DWZ10] ———, *Quivers with potentials and their representations II: Applications to cluster algebras*, J. Amer. Math. Soc. **23** (2010), no. 3, 749–790, [arXiv:0904.0676v2](#).
- [Din10] Ming Ding, *On quantum cluster algebras of finite type*, [arXiv:1011.1606](#).
- [DX11] Ming Ding and Fan Xu, *A quantum analogue of generic bases for affine cluster algebras*, [arXiv:1105.2421v1](#).
- [Efi11] Alexander Ivanovich Efimov, *Quantum cluster variables via vanishing cycles*, 2011, [arXiv:1112.3601v2](#).
- [FZ02] Sergey Fomin and Andrei Zelevinsky, *Cluster algebras. I. Foundations*, J. Amer. Math. Soc. **15** (2002), no. 2, 497–529 (electronic), [arXiv:math/0104151v1](#).
- [FZ07] ———, *Cluster algebras IV: Coefficients*, Compositio Mathematica **143** (2007), 112–164, [arXiv:math/0602259v3](#).
- [GLS11] Christof Geiß, Bernard Leclerc, and Jan Schröer, *Kac-Moody groups and cluster algebras*, Advances in Mathematics **228** (2011), no. 1, 329–433, [arXiv:1001.3545v2](#).

- [GLS12] ———, *Generic bases for cluster algebras and the Chamber Ansatz*, J. Amer. Math. Soc. **25** (2012), no. 1, 21–76, [arXiv:1004.2781v3](#).
- [HV08] Tamás Hausel and Fernando Rodriguez Villegas, *Mixed Hodge polynomials of character varieties*, Invent. Math. **174** (2008), no. 3, 552–624, with an appendix by Nicholas M. Katz, [arXiv:math/0612668v3](#).
- [HL10] David Hernandez and Bernard Leclerc, *Cluster algebras and quantum affine algebras*, Duke Math. J. **154** (2010), no. 2, 265–341, [arXiv:0903.1452](#).
- [HL11] ———, *Quantum Grothendieck rings and derived Hall algebras*, 2011, [arXiv:1109.0862v1](#).
- [IY08] Osamu Iyama and Yuji Yoshino, *Mutations in triangulated categories and rigid Cohen-Macaulay modules*, Inv. Math. **172** (2008), 117–168.
- [Kel12] Bernhard Keller, *Cluster algebras and derived categories*, 2012, [arXiv:1202.4161](#).
- [Kim12] Yoshiyuki Kimura, *Quantum unipotent subgroup and dual canonical basis*, Kyoto J. Math. **52** (2012), no. 2, 277–331, [arXiv:1010.4242](#), [doi:10.1215/21562261-1550976](#).
- [KQ12] Yoshiyuki Kimura and Fan Qin, *Graded quiver varieties, quantum cluster algebras and dual canonical basis*, 2012, [arXiv:1205.2066](#).
- [KS08] Maxim Kontsevich and Yan Soibelman, *Stability structures, Donaldson-Thomas invariants and cluster transformations*, 2008, [arXiv:0811.2435v1](#).
- [KY11] Bernhard Keller and Dong Yang, *Derived equivalences from mutations of quivers with potential*, Adv. Math. **226** (2011), no. 3, 2118–2168, [arXiv:0906.0761v3](#).
- [Lam11a] Philipp Lampe, *A quantum cluster algebra of Kronecker type and the dual canonical basis*, Int. Math. Res. Not. IMRN (2011), no. 13, 2970–3005, [arXiv:1002.2762v2](#), [doi:10.1093/imrn/rnq162](#).
- [Lam11b] ———, *Quantum cluster algebras of type A and the dual canonical basis*, 2011, [arXiv:1101.0580v1](#).
- [Lus90] G. Lusztig, *Canonical bases arising from quantized enveloping algebras*, J. Amer. Math. Soc. **3** (1990), no. 2, 447–498.
- [Lus00] ———, *Semicanonical bases arising from enveloping algebras*, Adv. Math. **151** (2000), no. 2, 129–139.
- [MSW11] Gregg Musiker, Ralf Schiffler, and Lauren Williams, *Bases of cluster algebras from surfaces*, 2011, [arXiv:1110.4364v2](#).
- [Nag10] Kentaro Nagao, *Donaldson-Thomas theory and cluster algebras*, 2010, [arXiv:1002.4884](#).
- [Nak94] Hiraku Nakajima, *Instantons on ALE spaces, quiver varieties, and Kac-Moody algebras*, Duke Math. J. **76** (1994), no. 2.
- [Nak98] ———, *Quiver varieties and Kac-Moody algebras*, Duke Math. J. **91** (1998), no. 3.
- [Nak01a] ———, *Quiver varieties and finite-dimensional representations of quantum affine algebras*, J. Amer. Math. Soc. **14** (2001), no. 1, 145–238 (electronic), [arXiv:math/9912158](#).
- [Nak01b] ———, *Quiver varieties and tensor products*, Invent. Math. **146** (2001), no. 2, 399–449, [arXiv:math.QA/0103008](#).
- [Nak04] ———, *Quiver varieties and  $t$ -analogs of  $q$ -characters of quantum affine algebras*, Ann. of Math. (2) **160** (2004), no. 3, 1057–1097, [arXiv:math/0105173v2](#).
- [Nak11] ———, *Quiver varieties and cluster algebras*, Kyoto J. Math. **51** (2011), no. 1, 71–126, [arXiv:0905.0002v5](#).
- [Pal08] Yann Palu, *Cluster characters for 2-Calabi-Yau triangulated categories*, Ann. Inst. Fourier (Grenoble) **58** (2008), no. 6, 2221–2248, [arXiv:math/0703540v2](#).
- [Pla10a] Pierre-Guy Plamondon, *Cluster algebras via cluster categories with infinite-dimensional morphism spaces*, 2010, [arXiv:1004.0830v1](#).
- [Pla10b] ———, *Cluster characters for cluster categories with infinite-dimensional morphism spaces*, 2010, [arXiv:1002.4956v2](#).
- [Pla11] ———, *Catégories amassées aux espaces de morphismes de dimension infinie, applications*, Ph.D. thesis, Université Paris Diderot - Paris 7, 2011, [http://people.math.jussieu.fr/~plamondon/plamondon\\_these.pdf](http://people.math.jussieu.fr/~plamondon/plamondon_these.pdf).

- [Qin10] Fan Qin, *Quantum cluster variables via Serre polynomials*, with an appendix by Bernhard Keller, to appear in *Journal für die reine und angewandte Mathematik* (Crelle's Journal), [arXiv:1004.4171](#), [doi:10.1515/CRELLE.2011.129](#).
- [Rei08] Markus Reineke, *Framed quiver moduli, cohomology, and quantum groups*, *J. Algebra* **320** (2008), no. 1, 94–115, [doi:10.1016/j.jalgebra.2008.01.025](#). MR 2417980 (2009d:16021)
- [Slo80] P. Slodowy, *Four lectures on simple groups and singularities*, Communications of the Mathematical Institute, Mathematical Institute, Rijksuniversitet Utrecht, 1980.
- [Tra09] Thao Tran, *F-polynomials in quantum cluster algebras*, 2009, [arXiv:0904.3291v1](#).
- [VV03] M. Varagnolo and E. Vasserot, *Perverse sheaves and quantum Grothendieck rings*, Studies in memory of Issai Schur (Chevaleret/Rehovot, 2000), *Progr. Math.*, vol. 210, Birkhäuser Boston, Boston, MA, 2003, pp. 345–365, [arXiv:math/0103182v3](#). MR MR1985732 (2004d:17023)

FAN QIN, BUREAU 7C01, UNIVERSITÉ PARIS DIDEROT - PARIS 7, INSTITUT DE MATHÉMATIQUES DE JUSSIEU, UMR 7586 DU CNRS, 175 RUE DU CHEVALERET, 75013, PARIS, FRANCE

*E-mail address:* [qinfan@math.jussieu.fr](mailto:qinfan@math.jussieu.fr)